



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/42		A1	(11) International Publication Number: WO 99/01543
			(43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/DK98/00286 (22) International Filing Date: 30 June 1998 (30.06.98) (30) Priority Data: 0812/97 4 July 1997 (04.07.97) DK 0846/97 11 July 1997 (11.07.97) DK 60/053,506 23 July 1997 (23.07.97) US (71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors: BJØRNVAD, Mads, Eskelund; Dr. Abildgaards Allé 8, 3th, DK-1955 Frederiksberg (DK). HATAKEYAMA, Mariko; 2257 Fox Glen Drive, Fairfield, CA 94533 (US). SCHÜLEIN, Martin; Wiedeweltsgade 51, DK-2100 Copenhagen Ø (DK). NIELSEN, Jack, Bech; Ole Olsens Allé 12, DK-2900 Hellerup (DK).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: ENDO- β -1,4-GLUCANASES FROM SACCHAROTHRIX			
(57) Abstract			
<p>An enzyme preparation comprising an enzyme having endo-β-1,4-glucanase activity obtainable from or endogeneous to a strain belonging to the genus <i>Saccharothrix</i> such as <i>Saccharothrix australiensis</i>, IFO 14444; an isolated polynucleotide (DNA) molecule encoding an enzyme or enzyme core (the catalytically active domain of the enzyme) exhibiting endo-β-1,4-glucanase activity selected from: (a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 676 to nucleotide 1470; (b) polynucleotide molecules that encode a polypeptide that is at least 80 % identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 226 to amino acid residue 490; and (c) degenerate nucleotide sequences of (a) or (b), the expressed endoglucanase and the enzyme preparation being useful in a detergent or fabric softener composition or in the textile industry for improving the properties of cellulosic fibres or fabric or for providing a stone-washed look of denim.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ENDO- β -1,4-GLUCANASES FROM *SACCHAROTHRIX*

The present invention relates to an endo- β -1,4-glucanase preparation derived from a *Saccharothrix* strain, an isolated polynucleotide molecule encoding an endo- β -1,4-glucanase, an isolated enzyme produced by recombinant techniques, and use of the preparation or the enzyme in the detergent, paper and pulp, oil drilling, oil extraction, wine and juice, food ingredients, animal feed and textile industries.

10 BACKGROUND OF THE INVENTION

Cellulose is a polymer of glucose linked by β -1,4-glucosidic bonds. Cellulose chains form numerous intra- and intermolecular hydrogen bonds, which result in the formation of insoluble cellulose microfibrils. Microbial hydrolysis of cellulose to glucose involves the following three major classes of cellulases: (i) endoglucanases (EC 3.2.1.4) which cleave β -1,4-glucosidic links randomly throughout cellulose molecules; (ii) cellobiohydrolases (EC 3.2.1.91) which digest cellulose from the nonreducing end, releasing cellobiose; and (iii) β -glucosidases (EC 3.2.1.21) which hydrolyse cellobiose and low-molecular-mass cellodextrins to release glucose. Cellulases are produced by many microorganisms and are often present in multiple forms. Recognition of the economic significance of the enzymatic degradation of cellulose has promoted an extensive search for microbial cellulases which can be used industrially. As a result, the enzymatic properties and the primary structures of a large number of cellulase have been investigated. On the basis of the results of a hydrophobic cluster analysis of the amino acid sequence of the catalytic domain, these cellulases have been placed into different families of glycosyl hydrolases; fungal and bacterial glycosyl hydrolases have been grouped into 35 families (Henrissat et. al. (1991), (1993)). Most cellulases consist of a cellulose-binding domain (CBD) and a catalytic domain (CAD) separated by a linker which may be rich in proline and hydroxy amino residues. Another classification of cellulases has been established on the basis

of the similarity of their CBDs (Gilkes et al. (1991)) giving five families of glycosyl hydrolases (I-V).

Cellulases are synthesized by a large number of microorganisms which include fungi, actinomycetes, myxobacteria and true bacteria but also by plants. Especially endo- β -1,4-glucanases of a wide variety of specificities have been identified. Many bacterial endoglucanases have been described (Henrissat (1993); Gilbert et al., (1993)).

According to the NCBI Taxonomy Browser (Internet address: <http://www.ncbi.nlm.nih.gov/Taxonomy/tax.html>), the taxonomic family *Pseudonocardiaceae* belongs to the order of *Actinomycetes* under the class of gram-positive bacteria (*Firmicutes*) and covers the genera *Actinokineospora*, *Actinopolyspora*, *Amycolata*, *Amycolatopsis*, *Kibdelosporangium*, *Pseudonocardia*, *Saccharomonospora*, *Saccharopolyspora*, *Saccharothrix*, and *Thermocrispum*. According to Taxonomy of bacteria and Archaea refers to the Ribosomal Database Project (B.L. Maidak et al. (1996) *Nucleic Acids Res.* 24:82-85), the Gram-Positive Phylum, High mol% G+C Subdivision comprises *Streptomyces* group (e.g. *Kitasatosporia setae*, *Streptomyces albogriseolus*, *Streptomyces lividans*, *Streptomyces cellulflavus*, *Streptomyces flavogriseus* and *Streptomyces griseus*), *Arthrobacter* group (e.g. *Cellulomonas fimi*, *Cellulomonas gelida*, *Cellulomonas flavigena* and *Cellulomonas uda*), and *Saccharopolyspora* group (e.g. *Thermomonospora curvata*, *Thermomonospora mesouviiformis*, *micromonospora inositola*, *Saccharothrix australiensis*, *Saccharothrix longispora*, *Saccharothrix mutabilis*, *Amycolatopsis mediterranei* and *Nocardiopsis albus*).

A very important industrial use of cellulolytic enzymes is the use for treatment of cellulosic textile or fabric, e.g. as ingredients in detergent compositions or fabric softener compositions, for bio-polishing of new fabric (garment finishing), and for obtaining a "stone-washed" look of cellulose-containing fabric, especially denim, and several methods for such treatment have been suggested, e.g. in GB-A-1 368 599, EP-A-0 307 564 and EP-A-0 435 876, WO 91/17243, WO 91/10732, WO 91/17244, PCT/DK95/000108 and PCT/DK95/00132. Another important industrial use of cellulolytic enzymes is the use for treatment

of paper pulp, e.g. for improving the drainage or for deinking of recycled paper.

There is an ever existing need for providing novel cellulase enzymes or enzyme preparations which may be used for applications where cellulase, preferably an endo- β -1,4-glucanase, activity is desirable.

The object of the present invention is to provide novel enzymes and enzyme compositions having substantial cellulolytic activity under slightly acidic to alkaline conditions and improved performance in paper pulp processing, textile treatment, laundry processes, extraction processes or in animal feed; preferably are such novel well-performing endoglucanases producible or produced by using recombinant techniques in high yields.

SUMMARY OF THE INVENTION

It has now been found that a variety of strains belonging to the genus *Saccharothrix* produces an enzyme having substantial cellulolytic activity, i.e. an endo- β -1,4-glucanase which is endogenous to *Saccharothrix*, and the inventors have succeeded in cloning and expressing a DNA sequence encoding such an enzyme.

Accordingly, in its first aspect the present invention relates to an enzyme preparation comprising an enzyme having endo- β -1,4-glucanase activity which is obtainable from or endogeneous to a strain belonging to the genus *Saccharothrix*, preferably belonging to a strain selected from the species *Saccharothrix australiensis*, *Saccharothrix texasensis*, *Saccharothrix waywayandensis*, *Saccharothrix cryophilis*, *Saccharothrix flava*, *Saccharothrix coeruleofusca*, *Saccharothrix longispora*, *Saccharothrix mutabilis* ssp. *capreolus*, *Saccharothrix aerocolonigenes*, *Saccharothrix mutabilis* ssp. *mutabilis*, *Saccharothrix syringae*, *Saccharothrix* sp., and in its second aspect the invention relates to an isolated polynucleotide molecule, preferably a DNA molecule, encoding an enzyme or enzyme core (ie the catalytically active domain of the enzyme) exhibiting endo- β -1,4-glucanase activity selected from the group consisting of (a) polynucleotide molecules comprising a nucleo-

tide sequence as shown in SEQ ID NO:1 from nucleotide 676 to nucleotide 1470, (b) polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 226 to amino acid residue 490, and (c) degenerate nucleotide sequences of (a) or (b); preferably a polynucleotide molecule capable of hybridizing to a denatured double-stranded DNA probe under medium stringency conditions, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 676-1470 of SEQ ID NO:1 and DNA probes comprising a subsequence of positions 676-1470 of SEQ ID NO:1 having a length of at least about 100 base pairs.

A plasmid pSJ1678 comprising a DNA sequence encoding the endoglucanase of the invention has been transformed into a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 17 March 1997 under the deposition number DSM 11476.

In its third, fourth and fifth aspect the invention provides an expression vector comprising a DNA segment which is eg a polynucleotide molecule of the invention; a cell comprising the DNA segment or the expression vector; and a method of producing an enzyme exhibiting cellulolytic activity, which method comprises culturing the cell under conditions permitting the production of the enzyme, and recovering the enzyme from the culture.

In yet another aspect the invention provides an isolated enzyme exhibiting cellulolytic activity, characterized in (i) being free from homologous impurities and (ii) the enzyme is produced by the method described above.

The invention further relates to an endo-beta-1,4-glucanase being either a polypeptide produced by *Saccharothrix australiensis*, IFO 14444, or a polypeptide comprising an amino acid sequence as shown in positions 226-490 of SEQ ID NO:2, or an analogue of any of these polypeptides which is at least 75%

homologous therewith, or is derived from any of these polypeptides by substitution, deletion or addition of one or several amino acids, or is immunologically reactive with a polyclonal antibody raised against any of these polypeptides in
5 purified form.

Further, the present invention relates to the use of such an enzyme or the enzyme preparation of the invention for industrial applications such as in a detergent composition or fabric softener composition intended for domestic or industrial
10 use in conventional laundry washing processes; in industrial cleaning processes; and in the textile industry for improving the properties of cellulosic fibres or fabric or for providing a stone-washed look of denim.

The invention also relates to an isolated substantially
15 pure biological culture of the *Escherichia coli* strain DSM 11476 harbouring a cellulase encoding DNA sequence (the cellulase encoding part of the DNA sequence cloned into plasmid pSJ1678 present in *Escherichia coli* DSM 11476 derived from a strain of the bacterial species *Saccharothrix australiensis*, or
20 any mutant of said *E.coli* strain.

DETAILED DESCRIPTION OF THE INVENTION

The enzyme or enzyme preparation of the invention is obtainable from or endogeneous to a strain belonging to the
25 genus *Saccharothrix*, preferably from the species selected from the group consisting of *Saccharothrix australiensis*, *Saccharothrix texasensis*, *Saccharothrix waywayandensis*, *Saccharothrix cryophilis*, *Saccharothrix flava*, *Saccharothrix coeruleofusca*, *Saccharothrix longispora*, *Saccharothrix mutabilis*
30 *ssp. capreolus*, *Saccharothrix aerocolonigenes*, *Saccharothrix mutabilis ssp. mutabilis*, *Saccharothrix syringae*, *Saccharothrix sp.* Examples of useful strains are *Saccharothrix australiensis*, IFO 14444, *Saccharothrix texasensis*, NRRL B-16134, *Saccharothrix waywayandensis*, NRRL B-16159, *Saccharothrix cryophilis*, NRRL B-
35 16238, *Saccharothrix sp.*, IFO 13785, *Saccharothrix flava*, ATCC 29533, *Saccharothrix coeruleofusca* (*Actinomadura coeruleofusca*), ATCC 35108 or DSM 43679, *Saccharothrix longispora* (*Actinomadura longispora*), ATCC 31109 or DSM 43749, *Saccharothrix mutabilis*

ssp. mutabilis, ATCC 31520, *Saccharothrix aerocolonigenes*, ATCC 23870, *Saccharothrix mutabilis ssp. capreolus*, ATCC 23892, *Saccharothrix mutabilis*, DSM 40225 or DSM 43853, *Saccharothrix syringae*, DSM 43886.

5 The enzyme and the enzyme preparation of the invention is active over a broad pH range, preferably active at a pH between about 4 and about 11, preferably between about 5.5 and about 10.

 In a preferred embodiment, the enzyme or the enzyme preparation of the invention belongs to family 6 of glycosyl
10 hydrolases (Henrissat et al., *op.cit.*).

 In the present context the term "expression vector" denotes a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional
15 segments may include promoter and terminator sequences, and may optionally include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, and the like. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both. The
20 expression vector of the invention may be any expression vector that is conveniently subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which the vector is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as
25 an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

30 The term "recombinant expressed" or "recombinantly expressed" used herein in connection with expression of a polypeptide or protein is defined according to the standard definition in the art. Recombinantly expression of a protein is generally performed by using an expression vector as described
35 immediately above.

 The term "isolated", when applied to a polynucleotide molecule, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or

unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated
5 DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example,
10 Dynan and Tijan, Nature 316:774-78, 1985). The term "an isolated polynucleotide" may alternatively be termed "a cloned polynucleotide".

When applied to a protein/polypeptide, the term "isolated" indicates that the protein is found in a condition other than
15 its native environment. In a preferred form, the isolated protein is substantially free of other proteins, particularly other homologous proteins (*i.e.* "homologous impurities" (see below)). It is preferred to provide the protein in a greater than 40% pure form, more preferably greater than 60% pure form.

20 Even more preferably it is preferred to provide the protein in a highly purified form, *i.e.*, greater than 80% pure, more preferably greater than 95% pure, and even more preferably greater than 99% pure, as determined by SDS-PAGE.

The term "isolated protein/polypeptide" may alternatively be
25 termed "purified protein/polypeptide".

The term "homologous impurities" means any impurity (*e.g.* another polypeptide than the polypeptide of the invention) which originate from the homologous cell where the polypeptide of the invention is originally obtained from.

30 The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide and/or polypeptide produced by the specific source, or by a cell in which a gene from the source have been inserted.

The term "operably linked", when referring to DNA segments,
35 denotes that the segments are arranged so that they function in concert for their intended purposes, *e.g.* transcription initiates in the promoter and proceeds through the coding segment to the terminator

The term "polynucleotide" denotes a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules.

The term "complements of polynucleotide molecules" denotes polynucleotide molecules having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "promoter" denotes a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger peptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

POLYNUCLEOTIDES:

Within preferred embodiments of the invention an isolated polynucleotide of the invention will hybridize to similar sized regions of SEQ ID No. 1, or a sequence complementary thereto, under at least medium stringency conditions.

In particular polynucleotides of the invention will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in positions 676-1470 of SEQ ID NO:1 or any probe comprising a subsequence of SEQ ID NO:1 having a length of at least about 100 base pairs under at least

medium stringency conditions, but preferably at high stringency conditions as described in detail below. Suitable experimental conditions for determining hybridization at medium, or high stringency between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking of the filter containing the DNA fragments or RNA to hybridize in 5 x SSC (Sodium chloride/Sodium citrate, Sambrook et al. 1989) for 10 min, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS and 100 μ g/ml of denatured sonicated salmon sperm DNA (Sambrook et al. 1989), followed by hybridization in the same solution containing a concentration of 10ng/ml of a random-primed (Feinberg, A. P. and Vogelstein, B. (1983) Anal. Biochem. 132:6-13), 32P-dCTP-labeled (specific activity higher than 1×10^9 cpm/ μ g) probe for 12 hours at ca. 45°C. The filter is then washed twice for 30 minutes in 2 x SSC, 0.5 % SDS at least 60°C (medium stringency), still more preferably at least 65°C (medium/high stringency), even more preferably at least 70°C (high stringency), and even more preferably at least 75°C (very high stringency).

Molecules to which the oligonucleotide probe hybridizes under these conditions are detected using a x-ray film.

As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for isolating DNA and RNA are well known in the art. DNA and RNA encoding genes of interest can be cloned in Gene Banks or DNA libraries by means of methods known in the art.

Polynucleotides encoding polypeptides having endogucanase activity of the invention are then identified and isolated by, for example, hybridization or PCR.

The present invention further provides counterpart polypeptides and polynucleotides from different bacterial strains (orthologs or paralogs). Of particular interest are endoglucanase polypeptides from gram-positive alkalophilic strains, including species of *Saccharothrix*.

Species homologues of a polypeptide with endoglucanase activity of the invention can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a DNA

sequence of the present invention can be cloned using chromosomal DNA obtained from a cell type that expresses the protein. Suitable sources of DNA can be identified by probing Northern blots with probes designed from the sequences disclosed
5 herein. A library is then prepared from chromosomal DNA of a positive cell line. A DNA sequence of the invention encoding an polypeptide having endoglucanase activity can then be isolated by a variety of methods, such as by probing with probes designed from the sequences disclosed in the present specification and
10 claims or with one or more sets of degenerate probes based on the disclosed sequences. A DNA sequence of the invention can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent 4,683,202), using primers designed from the sequences disclosed herein. Within an additional method, the DNA
15 library can be used to transform or transfect host cells, and expression of the DNA of interest can be detected with an antibody (monoclonal or polyclonal) raised against the endoglucanase cloned from *S.australienses*, IFO 14444, expressed and purified as described in Materials and Methods and Examples
20 1 and 3, or by an activity test relating to a polypeptide having endoglucanase activity.

The endoglucanase encoding part of the DNA sequence cloned into plasmid pSJ1678 present in *Escherichia coli* DSM 11476 and/or an analogue DNA sequence of the invention may be cloned
25 from a strain of the bacterial species *Saccharothrix australiensis*, preferably the strain IFO 14444, producing the enzyme with endoglucanase activity, or another or related organism as described herein.

Alternatively, the analogous sequence may be constructed on
30 the basis of the DNA sequence obtainable from the plasmid present in *Escherichia coli* DSM 11476 (which is believed to be identical to the attached SEQ ID NO:1), e.g be a sub-sequence thereof, and/or by introduction of nucleotide substitutions which do not give rise to another amino acid sequence of the
35 mannanase encoded by the DNA sequence, but which corresponds to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions which

may give rise to a different amino acid sequence (i.e. a variant of the mannan degrading enzyme of the invention).

POLYPEPTIDES:

5 The sequence of amino acids nos. 226-490 of SEQ ID NO: 2 is a mature endoglucanase sequence of the catalytic active domain.

 The present invention also provides endoglucanase polypeptides that are substantially homologous to the polypeptide of SEQ ID NO:2 and species homologs (paralogs or
10 orthologs) thereof. The term "substantially homologous" is used herein to denote polypeptides having 75%, preferably at least 80%, more preferably at least 85%, and even more preferably at least 90%, sequence identity to the sequence shown in amino acids nos. 226-490 of SEQ ID NO:2 or their orthologs or
15 paralogs. Such polypeptides will more preferably be at least 95% identical, and most preferably 98% or more identical to the sequence shown in amino acids nos. 226-490 of SEQ ID NO:2 or its orthologs or paralogs. Percent sequence identity is determined by conventional methods, by means of computer programs known in
20 the art such as GAP provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) as disclosed in Needleman, S.B. and Wunsch, C.D., (1970), Journal of Molecular Biology, 48, 443-453, which is
25 hereby incorporated by reference in its entirety. GAP is used with the following settings for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

 Sequence identity of polynucleotide molecules is determined by similar methods using GAP with the following settings for DNA
30 sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3.

 Substantially homologous proteins and polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor nature, that is conservative amino acid substitutions (see Table
35 2) and other substitutions that do not significantly affect the folding or activity of the protein or polypeptide; small deletions, typically of one to about 30 amino acids; and small

amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, a small linker peptide of up to about 20-25 residues, or a small extension that facilitates purification (an affinity tag), such as a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al., Methods Enzymol. 198:3, 1991. See, in general Ford et al., Protein Expression and Purification 2: 95-107, 1991, which is incorporated herein by reference. DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA).

However, even though the changes described above preferably are of a minor nature, such changes may also be of a larger nature such as fusion of larger polypeptides of up to 300 amino acids or more both as amino- or carboxyl-terminal extensions to a polypeptide of the invention having endoglucanase activity.

Table 1

Conservative amino acid substitutions

20	Basic:	arginine
		lysine
		histidine
	Acidic:	glutamic acid
		aspartic acid
25	Polar:	glutamine
		asparagine
	Hydrophobic:	leucine
		isoleucine
		valine
30	Aromatic:	phenylalanine
		tryptophan
		tyrosine
35	Small:	glycine
		alanine
		serine
		threonine
		methionine

In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline and α-methyl serine) may be substituted for amino acid residues of a polypeptide according to the invention. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, or preferably, are commercially available, and include pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Essential amino acids in the endoglucanase polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244: 1081-1085, 1989). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity (i.e. mannanase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699-4708, 1996. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992. The identities of essential amino acids can also be inferred from analysis of homologies with polypeptides which are related to a polypeptide according to the invention.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis, recombination and/or shuffling followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53-57, 1988),

Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-2156, 1989), WO95/17413, or WO 95/22625. Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, or recombination/shuffling of different mutations
5 (WO95/17413, WO95/22625), followed by selecting for functional a polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem. 30:10832-10837, 1991; Ladner et
10 al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988).

Mutagenesis/shuffling methods as disclosed above can be combined with high-throughput, automated screening methods to de-
15 tect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of
20 interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in the art can identify and/or prepare a variety of polypeptides that are substantially homologous to residues 226 to 490 of SEQ
25 ID NO: 2 and retain the endoglucanase activity of the wild-type protein.

The endoglucanase enzyme of the invention may, in addition to the enzyme core comprising the catalytically domain, also comprise a cellulose binding domain (CBD), the cellulose binding
30 domain and enzyme core (the catalytically active domain) of the enzyme being operably linked. The cellulose binding domain (CBD) may exist as an integral part the encoded enzyme, or a CBD from another origin may be introduced into the endoglucanase thus creating an enzyme hybride. In this context, the term
35 "cellulose-binding domain" is intended to be understood as defined by Peter Tomme et al. "Cellulose-Binding Domains: Classification and Properties" in "Enzymatic Degradation of Insoluble Carbohydrates", John N. Saddler and Michael H. Penner

(Eds.), ACS Symposium Series, No. 618, 1996. This definition classifies more than 120 cellulose-binding domains into 10 families (I-X), and demonstrates that CBDs are found in various enzymes such as cellulases, xylanases, mannanases, 5 arabinofuranosidases, acetyl esterases and chitinases. CBDs have also been found in algae, e.g. the red alga *Porphyra purpurea* as a non-hydrolytic polysaccharide-binding protein, see Tomme et al., *op.cit.* However, most of the CBDs are from cellulases and xylanases, CBDs are found at the N and C termini of proteins or 10 are internal. Enzyme hybrids are known in the art, see e.g. WO 90/00609 and WO 95/16782, and may be prepared by transforming into a host cell a DNA construct comprising at least a fragment of DNA encoding the cellulose-binding domain ligated, with or without a linker, to a DNA sequence encoding the endoglucanase 15 and growing the host cell to express the fused gene. Enzyme hybrids may be described by the following formula:

$$\text{CBD} - \text{MR} - \text{X}$$

wherein CBD is the N-terminal or the C-terminal region of an amino acid sequence corresponding to at least the cellulose- 20 binding domain; MR is the middle region (the linker), and may be a bond, or a short linking group preferably of from about 2 to about 100 carbon atoms, more preferably of from 2 to 40 carbon atoms; or is preferably from about 2 to to about 100 amino acids, more preferably of from 2 to 40 amino acids; and X is an 25 N-terminal or C-terminal region of a polypeptide encoded by the first or second DNA sequence of the invention.

In a preferred embodiment, the isolated polynucleotide molecule of the invention further comprises a partial DNA sequence encoding a cellulose binding domain (CBD). An example 30 of such a partial DNA sequence is the sequence corresponding to the nucleotides in positions 60-435 of SEQ ID NO:1 or the CBD encoding part of the DNA sequence cloned into the plasmid pSJ1678 present in *Escherichia coli*, DSM 11476. The isolated polynucleotide molecule of the invention may comprise a further 35 partial nucleotide sequence encoding a linking region, the linking region operably linking the cellulose binding domain (CBD) and the catalytically active domain (CAD) of the enzyme encoded by the nucleotide sequence comprised by the isolated

polynucleotide molecule. Preferably, the linking region consists of from about 2 amino acid residues to about 120 amino acid residues, especially 10-80 amino acid residues.

The polynucleotide molecules of the invention including
5 the partial DNA sequences described above can be cloned from the strain *Escherichia coli* DSM 11476 using standard methods e.g. as described by Sambrook et al., (1989), Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Lab.; Cold Spring Harbor, NY.

10 The polynucleotide molecules of the invention and partial DNA sequences thereof can also be cloned by any general method involving

- cloning, in suitable vectors, a DNA library from any organism, preferably a prokaryote, more preferably from a
15 bacterium, more preferably from a gram positive bacterium, especially from a strain belonging to the class *Actinomycetes*, more especially from the family *Pseudonocardiaceae*, for example from the genus *Saccharothrix*, especially from *Saccharothrix australiensis*, expected to produce the endo- β -1,4-glucanase of
20 interest,

- transforming suitable host cells with said vectors,
- culturing the host cells under suitable conditions to express any enzyme of interest encoded by a clone in the DNA library,
- 25 - screening for positive clones by determining any cellulolytic activity of the enzyme produced by such clones, and
- isolating the enzyme encoding DNA from such clones.

Alternatively, the DNA encoding a cellulase of the
30 invention may, in accordance with well-known procedures, conveniently be cloned from a suitable source, such as any of the below mentioned organisms, by use of synthetic oligonucleotide probes prepared on the basis of the DNA sequence obtainable from the plasmid present in *Escherichia coli* DSM
35 11476.

How to use a sequence of the invention to get other related sequences: The disclosed sequence information herein relating to a polynucleotide sequence encoding an endo-beta-1,4-

glucanase of the invention can be used as a tool to identify other homologous mannanases. For instance, polymerase chain reaction (PCR) can be used to amplify sequences encoding other homologous mannanases from a variety of microbial sources, in particular of different *Saccharothrix* species.

Immunological cross-reactivity

Polyclonal antibodies, especially monospecific polyclonal antibodies, to be used in determining immunological cross-reactivity may be prepared by use of a purified cellulolytic enzyme. More specifically, antiserum against the endoglucanase of the invention may be raised by immunizing rabbits (or other rodents) according to the procedure described by N. Axelsen et al. in: A Manual of Quantitative Immuno-electrophoresis, Blackwell Scientific Publications, 1973, Chapter 23, or A. Johnstone and R. Thorpe, Immunochemistry in Practice, Blackwell Scientific Publications, 1982 (more specifically p. 27-31). Purified immunoglobulins may be obtained from the antisera, for example by salt precipitation $((\text{NH}_4)_2 \text{SO}_4)$, followed by dialysis and ion exchange chromatography, e.g. on DEAE-Sephadex. Immunochemical characterization of proteins may be done either by Ouchterlony double-diffusion analysis (O. Ouchterlony in: Handbook of Experimental Immunology (D.M. Weir, Ed.), Blackwell Scientific Publications, 1967, pp. 655-706), by crossed immuno-electrophoresis (N. Axelsen et al., supra, Chapters 3 and 4), or by rocket immuno-electrophoresis (N. Axelsen et al., Chapter 2).

Microbial Sources

For the purpose of the present invention the term "obtained from" or "obtainable from" as used herein in connection with a specific source, means that the enzyme is produced or can be produced by the specific source, or by a cell in which a gene from the source have been inserted.

It is at present contemplated that the cellulase of the invention may be obtained from a gram positive bacterium belonging to the family Pseudonocardiaceae, in particular a

strain of the genus *Saccharothrix*, in particular a strain of *Saccharothrix australiensis*.

In a preferred embodiment, the cellulase of the invention is obtained from the strain *Saccharothrix australiensis*, IFO 5 14444. It is at present contemplated that a DNA sequence encoding an enzyme homologous to the enzyme of the invention may be obtained from other strains belonging to the genus *Saccharothrix*.

An isolate of a strain of *Saccharothrix australiensis* 10 from which an endo- β -1,4-glucanase of the invention can be derived is publicly available from Institute for Fermentation (IFO), 17-85 Juso-honmachi 2-chome, Yodogawa-ku, Osaka 532, Japan, under the deposition number IFO 14444.

Further, the plasmid pSJ1678 comprising the DNA sequence 15 encoding the endoglucanase of the invention has been transformed into a strain of the *Escherichia coli* and deposited under the deposition number DSM 11476.

Recombinant expression vectors

20 A recombinant vector comprising a DNA construct encoding the enzyme of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously 25 replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome in part or in its entirety 30 and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector in which the DNA sequence encoding the enzyme of the invention is operably linked to additional segments required for transcription of the 35 DNA. In general, the expression vector is derived from plasmid or viral DNA, or may contain elements of both. The term, "operably linked" indicates that the segments are arranged so that they function in concert for their intended purposes, e.g.

transcription initiates in a promoter and proceeds through the DNA sequence coding for the enzyme.

The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be
5 derived from genes encoding proteins either homologous or heterologous to the host cell.

Examples of suitable promoters for use in bacterial host cells include the promoter of the *Bacillus stearothermophilus* maltogenic amylase gene, the *Bacillus licheniformis* alpha-
10 amylase gene, the *Bacillus amyloliquefaciens* alpha-amylase gene, the *Bacillus subtilis* alkaline protease gen, or the *Bacillus pumilus* xylosidase gene, or the phage Lambda P_R or P_L promoters or the *E. coli* lac, trp or tac promoters.

The DNA sequence encoding the enzyme of the invention may
15 also, if necessary, be operably connected to a suitable terminator.

The recombinant vector of the invention may further comprise a DNA sequence enabling the vector to replicate in the host cell in question.

20 The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, or a gene encoding resistance to e.g. antibiotics like kanamycin, chloramphenicol, erythromycin, tetracycline, spectinomycine, or the like, or resistance to heavy metals or
25 herbicides.

To direct an enzyme of the present invention into the secretory pathway of the host cells, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) may be provided in the recombinant vector. The
30 secretory signal sequence is joined to the DNA sequence encoding the enzyme in the correct reading frame. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the enzyme. The secretory signal sequence may be that normally associated with the enzyme or may be from a gene
35 encoding another secreted protein.

The procedures used to ligate the DNA sequences coding for the present enzyme, the promoter and optionally the terminator and/or secretory signal sequence, respectively, or to assemble

these sequences by suitable PCR amplification schemes, and to insert them into suitable vectors containing the information necessary for replication or integration, are well known to persons skilled in the art (cf., for instance, Sambrook et al.,
5 op.cit.).

Host cells

The cloned DNA molecule introduced into the host cell may be either homologous or heterologous to the host in question. If
10 homologous to the host cell, i.e. produced by the host cell in nature, it will typically be operably connected to another promoter sequence or, if applicable, another secretory signal sequence and/or terminator sequence than in its natural environment. The term "homologous" is intended to include a DNA
15 sequence encoding an enzyme native to the host organism in question. The term "heterologous" is intended to include a DNA sequence not expressed by the host cell in nature. Thus, the DNA sequence may be from another organism, or it may be a synthetic sequence.

20 The host cell into which the cloned DNA molecule or the recombinant vector of the invention is introduced may be any cell which is capable of producing the desired enzyme and includes bacteria, yeast, fungi and higher eukaryotic cells.

Examples of bacterial host cells which on cultivation are
25 capable of producing the enzyme of the invention may be a gram-positive bacteria such as a strain of *Bacillus*, in particular *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus*
30 *megatherium*, *Bacillus stearothermophilus*, *Bacillus subtilis* and *Bacillus thuringiensis*, a strain of *Lactobacillus*, a strain of *Streptococcus*, a strain of *Streptomyces*, in particular *Streptomyces lividans* and *Streptomyces murinus*, or the host cell may be a gram-negative bacteria such as a strain of *Escherichia*
35 *coli*.

The transformation of the bacteria may be effected by protoplast transformation, electroporation, conjugation, or by

using competent cells in a manner known per se (cf. e.g. Sambrook et al., supra).

When expressing the enzyme in a bacteria such as *Escherichia coli*, the enzyme may be retained in the cytoplasm, typically as insoluble granules (known as inclusion bodies), or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed and the granules are recovered and denatured after which the enzyme is refolded by diluting the denaturing agent. In the latter case, the enzyme may be recovered from the periplasmic space by disrupting the cells, e.g. by sonication or osmotic shock, to release the contents of the periplasmic space and recovering the enzyme.

When expressing the enzyme in a gram-positive bacteria such as a strain of *Bacillus* or a strain of *Streptomyces*, the enzyme may be retained in the cytoplasm, or may be directed to the extracellular medium by a bacterial secretion sequence.

Examples of a fungal host cell which on cultivation are capable of producing the enzyme of the invention is e.g. a strain of *Aspergillus* or *Fusarium*, in particular *Aspergillus awamori*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, and *Fusarium oxysporum*, and a strain of *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma reesei* and *Trichoderma viride*.

Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known per se. The use of a strain of *Aspergillus* as a host cell is described in EP 238 023 (Novo Nordisk A/S), the contents of which are hereby incorporated by reference.

Examples of a host cell of yeast origin which on cultivation are capable of producing the enzyme of the invention is e.g. a strain of *Hansenula* sp., a strain of *Kluyveromyces* sp., in particular *Kluyveromyces lactis* and *Kluyveromyces marcianus*, a strain of *Pichia* sp., a strain of *Saccharomyces*, in particular *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces kluyveri* and *Saccharomyces uvarum*, a strain of *Schizosaccharomyces* sp., in particular

Schizosaccharomyces pombe, and a strain of *Yarrowia* sp., in particular *Yarrowia lipolytica*.

Examples of a host cell of plant origin which on cultivation are capable of producing the enzyme of the invention
5 is e.g. a plant cell of *Solanum tuberosum* or *Nicotiana tabacum*.

Method of producing a cellulolytic enzyme

The present invention provides a method of producing an isolated enzyme according to the invention, wherein a suitable
10 host cell, which has been transformed with a DNA sequence encoding the enzyme, is cultured under conditions permitting the production of the enzyme, and the resulting enzyme is recovered from the culture.

As defined herein, an isolated polypeptide (e.g. an
15 enzyme) is a polypeptide which is essentially free of other polypeptides, e.g., at least about 20% pure, preferably at least about 40% pure, more preferably about 60% pure, even more preferably about 80% pure, most preferably about 90% pure, and even most preferably about 95% pure, as determined by SDS-PAGE.

20 The term "isolated polypeptide" may alternatively be termed "purified polypeptide".

When an expression vector comprising a DNA sequence encoding the enzyme is transformed into a heterologous host cell it is possible to enable heterologous recombinant production of
25 the enzyme of the invention.

Thereby it is possible to make a highly purified or monocomponent cellulolytic composition, characterized in being free from homologous impurities.

In this context homologous impurities means any impurities
30 (e.g. other polypeptides than the enzyme of the invention) which originate from the homologous cell where the enzyme of the invention is originally obtained from.

In the present invention the homologous host cell may be a strain of *Saccharothrix australiensis*.

35 The medium used to culture the transformed host cells may be any conventional medium suitable for growing the host cells in question. The expressed cellulolytic enzyme may conveniently be secreted into the culture medium and may be recovered

therefrom by well-known procedures including separating the cells from the medium by centrifugation or filtration, precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by chromatographic
5 procedures such as ion exchange chromatography, affinity chromatography, or the like.

Enzyme compositions

In a still further aspect, the present invention relates
10 to an enzyme composition comprising an enzyme exhibiting cellulolytic activity as described above.

The enzyme composition of the invention may, in addition to the cellulase of the invention, comprise one or more other enzyme types, for instance hemicellulase such as xylanase and
15 mannanase, other cellulase or endo- β -1,4-glucanase components, chitinase, lipase, esterase, pectinase, cutinase, phytase, oxidoreductase (peroxidase, haloperoxidase, oxidase, laccase), protease, amylase, reductase, phenoloxidase, ligninase, pullulanase, pectate lyase, xyloglucanase, pectin acetyl esterase, polygalacturonase, rhamnogalacturonase, pectin lyase, pectin methylesterase, cellobiohydrolase, transglutaminase; or mixtures thereof.

The enzyme composition may be prepared in accordance with methods known in the art and may be in the form of a liquid or a
25 dry composition. For instance, the enzyme composition may be in the form of a granulate or a microgranulate. The enzyme to be included in the composition may be stabilized in accordance with methods known in the art.

Endoglucanases have potential uses in a lot of different
30 industries and applications. Examples are given below of preferred uses of the enzyme composition of the invention. The dosage of the enzyme composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

35 The enzyme composition according to the invention may be useful for at least one of the following purposes

Uses

During washing and wearing, dyestuff from dyed fabrics or garment will conventionally bleed from the fabric which then looks faded and worn. Removal of surface fibers from the fabric
5 will partly restore the original colours and looks of the fabric. By the term "colour clarification", as used herein, is meant the partly restoration of the initial colours of fabric or garment throughout multiple washing cycles.

The term "de-pilling" denotes removing of pills from the
10 fabric surface.

The term "soaking liquor" denotes an aqueous liquor in which laundry may be immersed prior to being subjected to a conventional washing process. The soaking liquor may contain one or more ingredients conventionally used in a washing or launder-
15 ing process.

The term "washing liquor" denotes an aqueous liquor in which laundry is subjected to a washing process, i.e. usually a combined chemical and mechanical action either manually or in a washing machine. Conventionally, the washing liquor is an
20 aqueous solution of a powder or liquid detergent composition.

The term "rinsing liquor" denotes an aqueous liquor in which laundry is immersed and treated, conventionally immediately after being subjected to a washing process, in order to rinse the laundry, i.e. essentially remove the detergent
25 solution from the laundry. The rinsing liquor may contain a fabric conditioning or softening composition.

The laundry subjected to the method of the present invention may be conventional washable laundry. Preferably, the major part of the laundry is sewn or unsewn fabrics, including knits,
30 wovens, denims, yarns, and toweling, made from cotton, cotton blends or natural or manmade cellulosics (e.g. originating from xylan-containing cellulose fibers such as from wood pulp) or blends thereof. Examples of blends are blends of cotton or rayon/viscose with one or more companion material such as wool,
35 synthetic fibers (e.g. polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyvinylidene chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers

(e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fibers, lyocell).

DETERGENT DISCLOSURE AND EXAMPLES

5 Surfactant system

The detergent compositions according to the present invention comprise a surfactant system, wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar
10 surfactants.

The surfactant is typically present at a level from 0.1% to 60% by weight.

The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or
15 gel compositions the surfactant is most preferably formulated in such a way that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

Preferred systems to be used according to the present invention comprise as a surfactant one or more of the nonionic
20 and/or anionic surfactants described herein.


Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of
25 alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment,
30 the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include IgepalTM CO-630, marketed by the GAF Corporation; and TritonTM X-45, X-114,
35 X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).

The condensation products of primary and secondary aliphatic alcohols with about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include TergitolTM 15-S-9 (The condensation product of C₁₁-C₁₅ linear alcohol with 9 moles ethylene oxide), TergitolTM 24-L-6 NMW (the condensation product of C₁₂-C₁₄ primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; NeodolTM 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), NeodolTM 23-3 (the condensation product of C₁₂-C₁₃ linear alcohol with 3.0 moles of ethylene oxide), NeodolTM 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), NeodolTM 45-5 (the condensation product of C₁₄-C₁₅ linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company, KyroTM EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol LA 050 (the condensation product of C₁₂-C₁₄ alcohol with 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant systems of the present invention are alkylpolysaccharides disclosed in US 4,565,647, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to

about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted
5 for the glucosyl moieties (optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-
10 positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula



15 wherein R^2 is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and
20 x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to
25 form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4-, and/or 6-position, preferably predominantly the 2-position.


The condensation products of ethylene oxide with a
30 hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight from about 1500 to about 1800 and will
35 exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the

polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially available
 5 PluronicTM surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and
 10 ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product
 15 contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available TetronicTM compounds, marketed by BASF.

20 Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethyleneoxide, alkylpolysaccharides, and
 25 mixtures hereof. Most preferred are C₈-C₁₄ alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C₈-C₁₈ alcohol ethoxylates (preferably C₁₀ avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.

Highly preferred nonionic surfactants are polyhydroxy fatty acid
 30 amide surfactants of the formula




wherein R¹ is H, or R¹ is C₁₋₄ hydrocarbyl, 2-hydroxyethyl, 2-
 35 hydroxypropyl or a mixture thereof, R² is C₅₋₃₁ hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof. Preferably, R¹ is methyl, R² is

straight C₁₁₋₁₅ alkyl or C₁₆₋₁₈ alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose or lactose, in a reductive amination reaction.

5 Highly preferred anionic surfactants include alkyl alkoxyated sulfate surfactants. Examples hereof are water soluble salts or acids of the formula RO(A)_mSO₃M wherein R is an unsubstituted C₁₀-C₂₄ alkyl or hydroxyalkyl group having a C₁₀-C₂₄ alkyl component, preferably a C₁₂-C₂₀ alkyl or hydro-xyalkyl, 10 more preferably C₁₂-C₁₈ alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), 15 ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and 20 dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C₁₂-C₁₈ alkyl polyethoxylate (1.0) sulfate (C₁₂-C₁₈E(1.0)M), C₁₂-C₁₈ alkyl polyethoxylate (2.25) sulfate (C₁₂-C₁₈(2.25)M, and C₁₂-C₁₈ alkyl 25 polyethoxylate (3.0) sulfate (C₁₂-C₁₈E(3.0)M), and C₁₂-C₁₈ alkyl polyethoxylate (4.0) sulfate (C₁₂-C₁₈E(4.0)M), wherein M is conveniently selected from sodium and potassium.

Suitable anionic surfactants to be used are alkyl ester sulfonate surfactants including linear esters of C₈-C₂₀ 30 carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO₃ according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

35 The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula:



5 wherein R³ is a C₈-C₂₀ hydrocarbyl, preferably an alkyl, or combination thereof, R⁴ is a C₁-C₆ hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable
 10 salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R³ is C₁₀-C₁₆ alkyl, and R⁴ is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates
 15 wherein R³ is C₁₀-C₁₆ alkyl.

Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula ROSO₃M wherein R preferably is a C₁₀-C₂₄ hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C₁₀-C₂₀ alkyl com-
 20 ponent, more preferably a C₁₂-C₁₈ alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and
 25 dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of C₁₂-C₁₆ are preferred for lower wash temperatures (e.g. below about 50°C) and C₁₆-C₁₈ alkyl chains
 30 are preferred for higher wash temperatures (e.g. above about 50°C).

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for
 35 example, sodium, potassium, ammonium, and substituted ammonium salts such as mono- di- and triethanolamine salts) of soap, C₈-C₂₂ primary or secondary alkanesulfonates, C₈-C₂₄ olefinsulfonates, sulfonated polycarboxylic acids prepared by

sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C₈-C₂₄ alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty
5 acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C₁₂-C₁₈
10 monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C₆-C₁₂ diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl
15 polyethoxy carboxylates such as those of the formula
$$RO(CH_2CH_2O)_k-CH_2COO-M^+$$
 wherein R is a C₈-C₂₂ alkyl, k is an integer from 1 to 10, and M is a soluble salt forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated
20 resin acids present in or derived from tall oil.


Alkylbenzene sulfonates are highly preferred. Especially preferred are linear (straight-chain) alkyl benzene sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18 carbon atoms.

25 Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in US 3,929,678, (Column 23, line 58 through Column 29, line 23, herein incorporated by reference).

30 When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

The laundry detergent compositions of the present
35 invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.


Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbonyl group. Examples of such cationic surfactants include the ammonium surfactants such as
 5 alkyltrimethylammonium halogenides, and those surfactants having the formula:



wherein R^2 is an alkyl or alkyl benzyl group having from about 8
 10 to about 18 carbon atoms in the alkyl chain, each R^3 is selected from the group consisting of $-CH_2CH_2-$, $-CH_2CH(CH_3)-$, $-CH_2CH(CH_2OH)-$, $-CH_2CH_2CH_2-$, and mixtures thereof; each R^4 is selected from the group consisting of C_1-C_4 alkyl, C_1-C_4 hydroxyalkyl, benzyl ring structures formed by joining the two
 15 R^4 groups, $-CH_2CHOHCHOHCOR^6CHOHCH_2OH$, wherein R^6 is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R^5 is the same as R^4 or is an alkyl chain, wherein the total number of carbon atoms or R^2 plus R^5 is not more than about 18; each y is from 0 to about 10, and
 20 the sum of the y values is from 0 to about 15; and X is any compatible anion.

Highly preferred cationic surfactants are the water soluble quaternary ammonium compounds useful in the present composition having the formula:

25




wherein R_1 is C_8-C_{16} alkyl, each of R_2 , R_3 and R_4 is independently C_1-C_4 alkyl, C_1-C_4 hydroxy alkyl, benzyl, and $-$
 30 $(C_2H_4O)_xH$ where x has a value from 2 to 5, and X is an anion. Not more than one of R_2 , R_3 or R_4 should be benzyl.

The preferred alkyl chain length for R_1 is $C_{12}-C_{15}$, particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived
 35 synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for R_2R_3 and R_4 are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

- coconut trimethyl ammonium chloride or bromide;
- coconut methyl dihydroxyethyl ammonium chloride or bromide;
- 5 decyl triethyl ammonium chloride;
- decyl dimethyl hydroxyethyl ammonium chloride or bromide;
- C₁₂₋₁₅ dimethyl hydroxyethyl ammonium chloride or bromide;
- coconut dimethyl hydroxyethyl ammonium chloride or bromide;
- myristyl trimethyl ammonium methyl sulphate;
- 10 lauryl dimethyl benzyl ammonium chloride or bromide;
- lauryl dimethyl (ethenoxy)₄ ammonium chloride or bromide;
- choline esters (compounds of formula (i) wherein R₁ is



di-alkyl imidazolines [compounds of formula (i)].

Other cationic surfactants useful herein are also described in US 4,228,044 and in EP 000 224.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 25%, preferably from about 1% to about 8% by weight of such cationic surfactants.

Ampholytic surfactants are also suitable for use in the laundry detergent compositions of the present invention. These surfactants can be broadly described as aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents contains at least about 8 carbon atoms, typically from about 8 to about 18 carbon atoms, and at least one contains an anionic water-solubilizing group, e.g. carboxy, sulfonate, sulfate. See US 3,929,678 (column 19, lines 18-35) for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about


15%, preferably from about 1% to about 10% by weight of such ampholytic surfactants.

Zwitterionic surfactants are also suitable for use in laundry detergent compositions. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. See US 3,929,678 (column 19, line 38 through column 22, line 48) for examples of zwitterionic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; watersoluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula:



wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is

from 0 to about 3: and each R⁵ is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R⁵ groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include C₁₀-C₁₈ alkyl dimethyl amine oxides and C₈-C₁₂ alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such semi-polar nonionic surfactants.

15 **Builder system**

The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetraacetate, metal ion sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also be used herein.

Suitable builders can be an inorganic ion exchange material, commonly an inorganic hydrated aluminosilicate material, more particularly a hydrated synthetic zeolite such as hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered silicate consisting of sodium silicate (Na₂Si₂O₅).

Suitable polycarboxylates containing one carboxy group include lactic acid, glycolic acid and ether derivatives thereof as disclosed in Belgian Patent Nos. 831,368, 821,369 and 821,370. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycollic acid,

tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates described in German Offenle-enschrift 2,446,686, and 2,446,487, US 3,935,257 and the sulfinyl carboxylates described in Belgian Patent No. 840,623.

- 5 Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in Netherlands
- 10 Application 7205873, and the oxypolycarboxylate materials such as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

- Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829,
- 15 1,1,2,2,-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in US 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,082,179,
- 20 while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

- Alicyclic and heterocyclic polycarboxylates include cyclopentane-cis,cis-cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-
- 25 tetracarboxylates, 2,5-tetrahydro-furan-cis, discarboxylates, 2,2,5,5,-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates and carboxymethyl derivatives of polyhydric alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid, pyromellitic acid and
- 30 the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxy-carboxylates containing up to three carboxy groups per molecule, more particularly citrates.

- 35 Preferred builder systems for use in the present compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered

silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

A suitable chelant for inclusion in the detergent compositions in accordance with the invention is ethylenediamine-N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline earth metal, ammonium, or substituted ammonium salts thereof, or mixtures thereof. Preferred EDDS compounds are the free acid form and the sodium or magnesium salt thereof. Examples of such preferred sodium salts of EDDS include Na_2EDDS and Na_4EDDS .

Examples of such preferred magnesium salts of EDDS include MgEDDS and Mg_2EDDS . The magnesium salts are the most preferred for inclusion in compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a water-soluble carboxylate chelating agent such as citric acid.

Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms.

Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about 40,000.

Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition. Preferred levels of builder for liquid detergents are from 5% to 30%.

Enzymes

Preferred detergent compositions, in addition to the enzyme preparation of the invention, comprise other enzyme(s) which provides cleaning performance and/or fabric care benefits.

Such enzymes include proteases, lipases, cutinases, amylases, cellulases, peroxidases, oxidases (e.g. laccases).

Proteases: Any protease suitable for use in alkaline solutions can be used. Suitable proteases include those of animal, 5 vegetable or microbial origin. Microbial origin is preferred. Chemically or genetically modified mutants are included. The protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived 10 from Bacillus, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO 89/06270.

15 Preferred commercially available protease enzymes include those sold under the trade names Alcalase, Savinase, Primase, Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal, Maxapem, Properase, Purafect and Purafect OXP by Genencor International, and those 20 sold under the tradename Opticlean and Optimase by Solvay Enzymes. Protease enzymes may be incorporated into the compositions in accordance with the invention at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by 25 weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Lipases: Any lipase suitable for use in alkaline solutions 30 can be used. Suitable lipases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included.

Examples of useful lipases include a Humicola lanuginosa lipase, e.g., as described in EP 258 068 and EP 305 216, a 35 Rhizomucor miehei lipase, e.g., as described in EP 238 023, a Candida lipase, such as a C. antarctica lipase, e.g., the C. antarctica lipase A or B described in EP 214 761, a Pseudomonas lipase such as a P. alcaligenes and P. pseudoalcaligenes lipase,

e.g., as described in EP 218 272, a P. cepacia lipase, e.g., as described in EP 331 376, a P. stutzeri lipase, e.g., as disclosed in GB 1,372,034, a P. fluorescens lipase, a Bacillus lipase, e.g., a B. subtilis lipase (Dartois et al., (1993),
5 Biochemica et Biophysica acta 1131, 253-260), a B. stearo-thermophilus lipase (JP 64/744992) and a B. pumilus lipase (WO 91/16422).

Furthermore, a number of cloned lipases may be useful, including the Penicillium camembertii lipase described by
10 Yamaguchi et al., (1991), Gene 103, 61-67), the Geotricum candidum lipase (Schimada, Y. et al., (1989), J. Biochem., 106, 383-388), and various Rhizopus lipases such as a R. delemar lipase (Hass, M.J et al., (1991), Gene 109, 117-113), a R. niveus lipase (Kugimiya et al., (1992), Biosci. Biotech.
15 Biochem. 56, 716-719) and a R. oryzae lipase.

Other types of lipolytic enzymes such as cutinases may also be useful, e.g., a cutinase derived from Pseudomonas mendocina as described in WO 88/09367, or a cutinase derived from Fusarium solani pisi (e.g. described in WO 90/09446).

20 Especially suitable lipases are lipases such as M1 LipaseTM, Luma fastTM and LipomaxTM (Genencor), LipolaseTM and Lipolase UltraTM (Novo Nordisk A/S), and Lipase P "Amano" (Amano Pharmaceutical Co. Ltd.).

The lipases are normally incorporated in the detergent
25 composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a
30 level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Amylases: Any amylase (a and/or b) suitable for use in alkaline solutions can be used. Suitable amylases include those of bacterial or fungal origin. Chemically or genetically mod-
35 ified mutants are included. Amylases include, for example, α -amylases obtained from a special strain of B. licheniformis, described in more detail in GB 1,296,839. Commercially available amylases are DuramylTM, TermamylTM, FungamylTM and BANTM

(available from Novo Nordisk A/S) and RapidaseTM and Maxamyl PTM (available from Genencor).

The amylases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Cellulases: Any cellulase suitable for use in alkaline solutions can be used. Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Suitable cellulases are disclosed in US 4,435,307 which discloses fungal cellulases produced from *Humicola insolens*, in WO 96/34108 and WO 96/34092 which disclose bacterial alkalophilic cellulases (BCE 103) from *Bacillus*, and in WO 94/21801, US 5,475,101 and US 5,419,778 which disclose EG III cellulases from *Trichoderma*. Especially suitable cellulases are the cellulases having colour care benefits. Examples of such cellulases are cellulases described in European patent application No. 0 495 257 and the endoglucanase of the present invention. Commercially available cellulases include CelluzymeTM and CarezymeTM produced by a strain of *Humicola insolens* (Novo Nordisk A/S), KAC-500(B)TM (Kao Corporation), and PuradaxTM (Genencor International).

Cellulases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Peroxidases/Oxidases : Peroxidase enzymes are used in combination with hydrogen peroxide or a source thereof (e.g. a percarbonate, perborate or persulfate). Oxidase enzymes are used in combination with oxygen. Both types of enzymes are used for

"solution bleaching", i.e. to prevent transfer of a textile dye from a dyed fabric to another fabric when said fabrics are washed together in a wash liquor, preferably together with an enhancing agent as described in e.g. WO 94/12621 and WO 95/01426. Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically or genetically modified mutants are included.

Peroxidase and/or oxidase enzymes are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Mixtures of the above mentioned enzymes are encompassed herein, in particular a mixture of a protease, an amylase, a lipase and/or a cellulase.

The enzyme of the invention, or any other enzyme incorporated in the detergent composition, is normally incorporated in the detergent composition at a level from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level from 0.01% to 0.2% of enzyme protein by weight of the composition.

Bleaching agents

Additional optional detergent ingredients that can be included in the detergent compositions of the present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%. In

general, bleaching compounds are optional added components in non-liquid formulations, e.g. granular detergents.

The bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including
5 oxygen bleaches as well as others known in the art.

The bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts
10 thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in US 4,483,781, US 740,446, EP 0 133 354 and US
15 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxy-caproic acid as described in US 4,634,551.

Another category of bleaching agents that can be used encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric
20 acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product, preferably 1-5% by weight.

The hydrogen peroxide releasing agents can be used in
25 combination with bleach activators such as tetra-acetylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, described in US 4,412,934), 3,5-trimethyl-hexanoyloxybenzenesulfonate (ISONOBS, described in EP 120 591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a
30 peracid as the active bleaching species, leading to improved bleaching effect. In addition, very suitable are the bleach activators C8(6-octanamido-caproyl) oxybenzene-sulfonate, C9(6-nonanamido caproyl) oxybenzenesulfonate and C10 (6-decanamido caproyl) oxybenzenesulfonate or mixtures thereof. Also suitable
35 activators are acylated citrate esters such as disclosed in European Patent Application No. 91870207.7.

Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen

bleaching compounds for use in cleaning compositions according to the invention are described in application USSN 08/136,626.

The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore)
5 which is capable of generation of hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in European Patent Application EP 0 537 381.

Bleaching agents other than oxygen bleaching agents are
10 also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminium phthalocyanines. These materials can be deposited upon the substrate during the washing process. Upon irradiation
15 with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated bleaching process are described in US 4,033,718. Typically,
20 detergent composition will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

Bleaching agents may also comprise a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds described in "Efficient manganese catalysts for low-temperature
25 bleaching", Nature 369, 1994, pp. 637-639.

Suds suppressors

Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures.
30 Silicones can generally be represented by alkylated polysiloxane materials, while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic silicas of various types. These materials can be incorporated as particulates, in which the suds suppressor is advantageously
35 releasably incorporated in a water-soluble or waterdispersible, substantially non surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed

in a liquid carrier and applied by spraying on to one or more of the other components.

A preferred silicone suds controlling agent is disclosed in US 3,933,672. Other particularly useful suds suppressors are
5 the self-emulsifying silicone suds suppressors, described in German Patent Application DTOS 2,646,126. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a
10 mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

Such suds suppressor system are described in European Patent Application EP 0 593 841.

15 Especially preferred silicone suds controlling agents are described in European Patent Application No. 92201649.8. Said compositions can comprise a silicone/ silica mixture in combination with fumed nonporous silica such as Aerosil^R.

The suds suppressors described above are normally employed
20 at levels of from 0.001% to 2% by weight of the composition, preferably from 0.01% to 1% by weight.

Other components

Other components used in detergent compositions may be
25 employed such as soil-suspending agents, soil-releasing agents, optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or nonencapsulated perfumes.

Especially suitable encapsulating materials are water
30 soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.

Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid esters of substituted dicarboxylic acids such as described in US
35 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulation materials include N-Lok manufactured by National Starch. The N-Lok

encapsulating material consists of a modified maize starch and glucose. The starch is modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.


Antiredeposition and soil suspension agents suitable
5 herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as
10 copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the
15 composition.

Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-
20 ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' - bis-(2,4-dianilino-s-tri-azin-6 ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-
25 2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-so-dium 4,4'bis(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylami-no)stilbene-2,2'disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3, - triazole-2''-sulphonate and 4,4'-bis(2-
30 sulphostyryl)biphenyl.

Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from
35 0.25% to 2.5% by weight. These polymers and the previously mentioned homo- or co-polymeric poly-carboxylate salts are valuable for improving whiteness maintenance, fabric ash

deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present invention are conventionally copolymers or terpolymers of
 5 terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in US 4,116,885 and 4,711,730 and EP 0 272 033. A particular preferred polymer in accordance with EP 0 272 033 has the formula:



where PEG is $-(\text{OC}_2\text{H}_4)_n-$, PO is $(\text{OC}_3\text{H}_6\text{O})$ and T is $(\text{pOOC}_6\text{H}_4\text{CO})$.

Also very useful are modified polyesters as random
 15 copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1,2-propanediol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or 1,2-propanediol. The
 20 target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be endcapped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of
 25 ethylene glycol and/or 1,2-propanediol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of 1,2-propanediol, about 10% by weight ethylene glycol, about 13% by
 30 weight of dimethyl sulfobenzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3.000. The polyesters and their method of preparation are described in detail in EP 311 342.

35 Softening agents

Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type.

Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400898 and in US 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP 0 011 340 and their
5 combination with mono C₁₂-C₁₄ quaternary ammonium salts are disclosed in EP-B-0 026 528 and di-long-chain amides as disclosed in EP 0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP 0 299 575 and 0
10 313 146.

Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the
15 water-insoluble tertiary amines or dilong chain amide materials are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%,
20 normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

25

Polymeric dye-transfer inhibiting agents

The detergent compositions according to the present invention may also comprise from 0.001% to 10%, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of
30 polymeric dye-transfer inhibiting agents. Said polymeric dye-transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability of complexing or adsorbing the
35 fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash.

Especially suitable polymeric dye-transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures
5 thereof.

Addition of such polymers also enhances the performance of the enzymes according to the invention.

The detergent composition according to the invention can be in liquid, paste, gels, bars or granular forms.

10 Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molecular weights of 1000 to
15 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-
20 forming coating materials suitable for application by fluid bed techniques are given in GB 1483591.

Granular compositions according to the present invention can also be in "compact form", i.e. they may have a relatively higher density than conventional granular detergents, i.e. form
25 550 to 950 g/l; in such case, the granular detergent compositions according to the present invention will contain a lower amount of "Inorganic filler salt", compared to conventional granular detergents; typical filler salts are alkaline earth metal salts of sulphates and chlorides, typically
30 sodium sulphate; "Compact" detergent typically comprise not more than 10% filler salt. The liquid compositions according to the present invention can also be in "concentrated form", in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to con-
35 ventional liquid detergents. Typically, the water content of the concentrated liquid detergent is less than 30%, more preferably less than 20%, most preferably less than 10% by weight of the detergent compositions.

The compositions of the invention may for example, be formulated as hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the pretreatment of stained fabrics, rinse
5 added fabric softener compositions, and compositions for use in general household hard surface cleaning operations and dishwashing operations.

The following examples are meant to exemplify compositions for the present invention, but are not necessarily meant to
10 limit or otherwise define the scope of the invention. In the detergent compositions, the abbreviated component identifications have the following meanings:

- LAS: Sodium linear C₁₂ alkyl benzene sulphonate
15 TAS: Sodium tallow alkyl sulphate
XYAS: Sodium C_{1X} - C_{1Y} alkyl sulfate
SS: Secondary soap surfactant of formula 2-butyl octanoic acid
25EY: A C₁₂ - C₁₅ predominantly linear primary alcohol
20 condensed with an average of Y moles of ethylene oxide
45EY: A C₁₄ - C₁₅ predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
XYEYZS: C_{1X} - C_{1Y} sodium alkyl sulfate condensed with an average of Z moles of ethylene oxide per mole
25 Nonionic: C₁₃ - C₁₅ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under the tradename Plurafax LF404 by BASF GmbH
CFAA: C₁₂ - C₁₄ alkyl N-methyl glucamide
30 TFAA: C₁₆ - C₁₈ alkyl N-methyl glucamide
Silicate: Amorphous Sodium Silicate (SiO₂:Na₂O ratio = 2.0)
NaSKS-6: Crystalline layered silicate of formula d-Na₂Si₂O₅
Carbonate: Anhydrous sodium carbonate
Phosphate: Sodium tripolyphosphate
35 MA/AA: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000

Polyacrylate: Polyacrylate homopolymer with an average molecular weight of 8,000 sold under the tradename PA30 by BASF GmbH

Zeolite A: Hydrated Sodium Aluminosilicate of formula
 5 $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$ having a primary particle size in the range from 1 to 10 micrometers

Citrate: Tri-sodium citrate dihydrate

Citric: Citric Acid

Perborate: Anhydrous sodium perborate monohydrate bleach,
 10 empirical formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$

PB4: Anhydrous sodium perborate tetrahydrate

Percarbonate: Anhydrous sodium percarbonate bleach of empirical formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$

TAED: Tetraacetyl ethylene diamine

15 CMC: Sodium carboxymethyl cellulose

DETPMP: Diethylene triamine penta (methylene phosphonic acid), marketed by Monsanto under the Tradename Dequest 2060

PVP: Polyvinylpyrrolidone polymer

EDDS: Ethylenediamine-N, N'-disuccinic acid, [S,S] isomer
 20 in the form of the sodium salt

Suds Suppressor: 25% paraffin wax Mpt 50°C, 17% hydrophobic silica, 58% paraffin oil

Granular Suds suppressor: 12% Silicone/silica, 18% stearyl alcohol, 70% starch in granular form

25 Sulphate: Anhydrous sodium sulphate

HMWPEO: High molecular weight polyethylene oxide

TAE 25: Tallow alcohol ethoxylate (25)

Detergent Example I

30

A granular fabric cleaning composition in accordance with the invention may be prepared as follows:

	Sodium linear C_{12} alkyl	6.5
35	benzene sulfonate	
	Sodium sulfate	15.0
	Zeolite A	26.0
	Sodium nitrilotriacetate	5.0

51

	Enzyme of the invention	0.1
	PVP	0.5
	TAED	3.0
	Boric acid	4.0
5	Perborate	18.0
	Phenol sulphonate	0.1
	Minors	Up to 100

Detergent Example II

10 A compact granular fabric cleaning composition (density 800 g/l) in accord with the invention may be prepared as follows:

	45AS	8.0
	25E3S	2.0
15	25E5	3.0
	25E3	3.0
	TFAA	2.5
	Zeolite A	17.0
	NaSKS-6	12.0
20	Citric acid	3.0
	Carbonate	7.0
	MA/AA	5.0
	CMC	0.4
	Enzyme of the invention	0.1
25	TAED	6.0
	Percarbonate	22.0
	EDDS	0.3
	Granular suds suppressor	3.5
	water/minors	Up to 100%

30

Detergent Example III

Granular fabric cleaning compositions in accordance with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows:

35	LAS	10.7	-
	TAS	2.4	-
	TFAA	-	4.0
	45AS	3.1	10.0

52

	45E7	4.0	-
	25E3S	-	3.0
	68E11	1.8	-
	25E5	-	8.0
5	Citrate	15.0	7.0
	Carbonate	-	10
	Citric acid	2.5	3.0
	Zeolite A	32.1	25.0
	Na-SKS-6	-	9.0
10	MA/AA	5.0	5.0
	DETPMP	0.2	0.8
	Enzyme of the invention	0.10	0.05
	Silicate	2.5	-
	Sulphate	5.2	3.0
15	PVP	0.5	-
	Poly (4-vinylpyridine)-N-Oxide/copolymer of vinyl-imidazole and vinyl-pyrrolidone	-	0.2
20	Perborate	1.0	-
	Phenol sulfonate	0.2	-
	Water/Minors	Up to 100%	

Detergent Example IV

25 Granular fabric cleaning compositions in accordance with the invention which provide "Softening through the wash" capability may be prepared as follows:

	45AS	-	10.0
	LAS	7.6	-
30	68AS	1.3	-
	45E7	4.0	-
	25E3	-	5.0
	Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
35	Citrate	5.0	3.0
	Na-SKS-6	-	11.0
	Zeolite A	15.0	15.0
	MA/AA	4.0	4.0

53

	DETPMP	0.4	0.4
	Perborate	15.0	-
	Percarbonate	-	15.0
	TAED	5.0	5.0
5	Smectite clay	10.0	10.0
	HMWPEO	-	0.1
	Enzyme of the invention	0.10	0.05
	Silicate	3.0	5.0
	Carbonate	10.0	10.0
10	Granular suds suppressor	1.0	4.0
	CMC	0.2	0.1
	Water/Minors	Up to 100%	

Detergent Example V

- 15 Heavy duty liquid fabric cleaning compositions in accordance with the invention may be prepared as follows:

		I	II
	LAS acid form	-	25.0
20	Citric acid	5.0	2.0
	25AS acid form	8.0	-
	25AE2S acid form	3.0	-
	25AE7	8.0	-
	CFAA	5	-
25	DETPMP	1.0	1.0
	Fatty acid	8	-
	Oleic acid	-	1.0
	Ethanol	4.0	6.0
	Propanediol	2.0	6.0
30	Enzyme of the invention	0.10	0.05
	Coco-alkyl dimethyl	-	3.0
	hydroxy ethyl ammonium		
	chloride		
	Smectite clay	-	5.0
35	PVP	2.0	-
	Water / Minors	Up to 100%	

Textile applications

In another embodiment, the present invention relates to use of the endoglucanase of the invention in the bio-polishing process. Bio-Polishing is a specific treatment of the yarn surface which improves fabric quality with respect to handle and appearance without loss of fabric wettability. The most important effects of Bio-Polishing can be characterized by less fuzz and pilling, increased gloss/luster, improved fabric handle, increased durable softness and altered water absorbency. Bio-Polishing usually takes place in the wet processing of the manufacture of knitted and woven fabrics. Wet processing comprises such steps as e.g. desizing, scouring, bleaching, washing, dying/printing and finishing. During each of these steps, the fabric is more or less subjected to mechanical action. In general, after the textiles have been knitted or woven, the fabric proceeds to a desizing stage, followed by a scouring stage, etc. Desizing is the act of removing size from textiles. Prior to weaving on mechanical looms, warp yarns are often coated with size starch or starch derivatives in order to increase their tensile strength. After weaving, the size coating must be removed before further processing the fabric in order to ensure a homogeneous and wash-proof result. It is known that in order to achieve the effects of Bio-Polishing, a combination of cellulytic and mechanical action is required. It is also known that "super-softness" is achievable when the treatment with a cellulase is combined with a conventional treatment with softening agents. It is contemplated that use of the endoglucanase of the invention for bio-polishing of cellulosic fabrics is advantageous, e.g. a more thorough polishing can be achieved. Bio-polishing may be obtained by applying the method described e.g. in WO 93/20278.

Stone-washing

It is known to provide a "stone-washed" look (localized abrasion of the colour) in dyed fabric, especially in denim fabric or jeans, either by washing the denim or jeans made from such fabric in the presence of pumice stones to provide the desired localized lightening of the colour of the fabric or by treating the fabric enzymatically, in particular with cellulytic

enzymes. The treatment with an endoglucanase of the present invention may be carried out either alone such as disclosed in US 4,832,864, together with a smaller amount of pumice than required in the traditional process, or together with perlite
5 such as disclosed in WO 95/09225.

MATERIALS AND METHODS

Organisms:

Saccharothrix australiensis, IFO 14444, comprises the
10 cellulase encoding DNA sequence of the invention.

Escherichia coli, DSM 11476, containing the plasmid comprising the DNA sequence encoding the cellulolytic enzyme of the invention, in the cloning vector pSJ1678.

Other strains:

E. coli strain: Cells of *E. coli* SJ2 (Diderichsen, B. et al. (1990)), were prepared for and transformed by electroporation using a Gene PulserTM electroporator from BIO-RAD as described by the supplier.

20

Plasmids:

pSJ1678 (see WO 94/19454 which is hereby incorporated by reference).

General molecular biology methods:

DNA manipulations and transformations were performed using standard methods of molecular biology (Sambrook et al, (1989); Ausubel et al., (1995); Harwood et al., (1990)).

Enzymes for DNA manipulations were used according to the
30 specifications of the suppliers.

Isolation of the DNA sequence encoding the cellulolytic enzyme of the invention:

The DNA sequence, comprising the DNA sequence shown in
35 SEQ ID No. 1, encoding the endoglucanase of the invention, can be obtained from the deposited organism *E. coli*, DSM 11476, by extraction of plasmid DNA by methods known in the art (Sambrook et al. (1989)).

Cloning of the *Saccharothrix australiensis* endo- β -1,4-glucanase gene

Genomic DNA preparation:

5 Strain *Saccharothrix australiensis*, IFO 14444, was propagated on TY-agar medium at 30 °C for 2-3 days. Cells were harvested by scraping them of the plates, and genomic DNA isolated by the method described by Pitcher et al. (1989). The only modification to the published method is the adding of
10 Mutanolysin (Catalogue no. M-9901, SIGMA, USA) of 30 ug/ml to the lysis buffer originally only containing lysozyme.

Genomic library construction:

Genomic DNA was partially digested with restriction enzyme
15 Sau3A, and size-fractionated by electrophoresis on a 0.7 % agarose gel. Fragments between 2 and 7 kb in size were isolated by electrophoresis onto DEAE-cellulose paper (Dretzen et al., (1981)).

Isolated DNA fragments were ligated to BamHI digested
20 pSJ1678 plasmid DNA, and the ligation mixture was used to transform *E. coli* SJ2.

Cells were plated on LB agar plates containing 0.1% CMC (Sodium-Carboxy-Methyl-Cellulose, Aqualon, France) and 9 μ g/ml Chloramphenicol and incubated overnight at 37°C.

25

Identification of positive clones by colony hybridization

A DNA library in *E. coli*, constructed as described above, was screened on LB agar plates containing 0.1% CMC (Sodium-Carboxy-Methyl-Cellulose, Aqualon, France) and 9 μ g/ml
30 Chloramphenicol and incubated overnight at 37°C. The transformants were subsequently replica plated onto the same type of plates, and these new plates were incubated 8 hours or overnight at 37°C.

The original plates were coloured using 1 mg/ml of Congo
35 Red (SIGMA, USA). The coloring was continued for half an hour with moderate orbital shaking, after which the plates were washed two times 15 minutes using 1 M NaCl.

Yellowish halos appeared at positions where cellulase positive clones were present, from the replica plates these cellulase positive clones were rescued and restreaked onto LB agar plates containing 0.1% CMC and 9 μ g/ml Chloramphenicol and 5 incubated overnight at 37°C.

Characterization of positive clones:

From the restreaking plates the endoglucanase positive clones were obtained as single colonies, and plasmids were ex-
10 tracted. Phenotypes were confirmed by retransformation of *E.coli* SJ2, and plasmids characterized by restriction digests.

The endoglucanase gene was characterized by DNA sequencing using the Taq deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labeled terminators and 5 pmol of the primer
15 #3507:

5'-GGC TTT TAA GCC GTC TGT ACG -3'.

In another reaction the nucleotide sequence was determined using the primer #8392:

5'-CTC ACG TTA AGG GAT TTT GGT CTG G -3'

20 Analysis of the sequence data was performed according to Devereux et al., (1984). The sequence corresponds to the DNA sequence shown in SEQ ID No 1. This first DNA sequences derived in this way was used to design new primers for further sequencing of the endoglucanase gene, and again new primers were
25 designed, used for sequencing and so forth.

Media

TY and LB agar (as described in Ausubel et al., (1995)).

Determination of cellulolytic activity

The cellulolytic activity of endoglucanase is determined relative to an analytical standard and may be expressed in the unit ECU.

- 5 Cellulolytic enzymes hydrolyse CMC, thereby decreasing the viscosity of the incubation mixture. The resulting reduction in viscosity may be determined by a vibration viscosimeter (e.g. MIVI 3000 from Sofraser, France).

Determination of the cellulolytic activity, measured in terms
10 of ECU, may be determined according to the analysis method AF 301.1 which is available from the Applicant upon request.

The ECU assay quantifies the amount of catalytic activity present in the sample by measuring the ability of the sample to reduce the viscosity of a solution of carboxy-methylcellulose
15 (CMC). The assay is carried out at 40°C, pH 7.5 using a relative enzyme standard for reducing the viscosity of the CMC substrate.

The following non-limiting examples illustrate the invention.

20 **EXAMPLE 1**

Cloning and expression of an endo- β -1,4-glucanase from *Saccharothrix australiensis*, IFO 14444

Preparation of genomic DNA from *Saccharothrix australiensis*, IFO 14444, cloning of the endoglucanase gene and
25 DNA sequencing was performed as described in Materials and Methods. One positive transformant isolated was DSM 11476, containing the plasmid pSJ1678 containing an insert of approximately 2.300 base-pairs. This insert was partially DNA sequenced, and revealed the presence of the sequence of an
30 endoglucanase-encoding gene. According to the classification of glycosyl hydrolases (Henrissat et al., (1993)) the endoglucanase belongs to the family 6 of glycosyl hydrolases.

The nucleotide sequence is designated SEQ ID NO 1.

The DNA corresponding to the endo- β -1,4-glucanase gene is
35 obtainable from the plasmid obtainable from the strain deposited as DSM 11476.

EXAMPLE 2**Endo- β -1,4-glucanase from various *Saccharothrix* strains**

The following strains were tested:

- Saccharothrix texasensis*, NRRL B-16134,
5 *Saccharothrix waywayandensis*, NRRL B-16159,
Saccharothrix cryophilis, NRRL B-16238,
Saccharothrix sp., IFO 13785,
Saccharothrix flava, ATCC 29533,
Saccharothrix coeruleofusca, ATCC 35108,
10 *Saccharothrix longispora*, ATCC 31109,
Saccharothrix mutabilis ssp. *mutabilis*, ATCC 31520,
Saccharothrix aerocolonigenes, ATCC 23870,
Saccharothrix mutabilis ssp. *capreolus*, ATCC 23892,
Saccharothrix syringae, DSM 43886.

15

A. Southern hybridization test of *Saccharothrix* strainsGenomic DNA preparation

Each *Saccharothrix* strain (see above) was propagated on TY-agar medium supplemented with 2% soluble starch at 25 °C for
20 3-4 days. Cells were harvested, and genomic DNA isolated by the method described by Pitcher et al. (Pitcher, D. G., Saunders, N. A., Owen, R. J. (1989). Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. Lett. Appl. Microbiol., 8, 151-156).

25 Hybridization conditions

Suitable conditions for determining hybridization between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking of the filter containing the DNA fragments or RNA to hybridize in 5 x SSC (standard saline citrate) for 10 min, and
30 prehybridization of the filter in a solution of 5 x SSC (Sambrook et al. 1989), 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS and 100 μ g/ml of denatured sonicated salmon sperm DNA (Sambrook et al. 1989), followed by hybridization in the same solution containing a random-primed (Feinberg, A. P.
35 and Vogelstein, B. (1983) *Anal. Biochem.* **132**:6-13), 32 P-dCTP-labeled (specific activity > 1 x 10⁹ cpm/ μ g) probe for 12 hours at ca. 45°C. The filter is then washed two times for 30 minutes in 2 x SSC, 0.5 % SDS at preferably at least 55°C, more

preferably at least 60°C, more preferably at least 65°C, even more preferably at least 70°C, especially at least 75°C.

Experimental

Probe:

- 5 2.5 kb fragment from pMB145 (*Saccharothrix australiensis*, IFO 14444).

Result: All strains hybridized to BamHI.

Accordingly, all strains produce endo- β -1,4-glucanase belonging to family 6.

10

B. CMC-Congo Red Assay of *Saccharothrix* strains (identification of positive clones by colony hybridization)

Method:

Strains were grown on wheat bran medium (w. bran 4%, y. ext.

- 15 0.1% and agar 1.5%) at 30°C for 4 days. Agar plugs were placed on CMC assay plates (basal: CMC 0.1%, agar 1.5%) at various pHs (pH 5, 7, 9, 10). LAS (linear alkyl sulfate) was added to the final conc. of 0.1%. AZCL-HE was used at pH 7. Incubation was carried out at 40°C, overnight. Then the plate was stained with
20 Congo Red and the diameter of clearing zones (mm) were measured.

Results (diameter in mm):

Source	AZCL-HE	pH 7	pH 9
IFO 14444	12	17	13
NRRL B-16134	10	17	15
NRRL B-16159	9	13	12
NRRL B-16238	-	9	-
IFO 13785	22	23	16
ATCC 29533	8	9	-
ATCC 35108	13	16	15
ATCC 31109	13	18	10
ATCC 31520	18	21	16
ATCC 23870	13	16	10
ATCC 23892	13	16	13
DSM 43886	12	16	13

EXAMPLE 3**Purification and characterization of an endo- β -1,4-glucanase from *Saccharothrix australiensis*, IFO 14444**

The plasmid of example 1 was inserted into *E. coli* JM 109. The *E. coli* strain was grown in super broth (pancreatic digest of casein 32 g/l, yeast extract 20 g/l and NaCl 5 gram/l adjusted to pH 7.0) at 37°C for one day. Cells were collected and
5 lysed by sonication. The supernatant was freeze dried. 24 gram was obtained from 13 l of fermentation with a total activity of 4600 ECU.

Purification:

The powder was diluted in ion exchanged water total of 175
10 ml. Using 50 gram of Avicel at 4°C, on a column 50 mM phosphate buffer pH 7.5 and washed using high salt (0.5 M NaCl in the same buffer). The enzyme eluted using ion exchanged pure water. Total 2550 ECU was obtained. The sample still contained some color and was further purified using ion exchange chromatography. A HPQ
15 column equilibrated with 50 mM Tris pH 7.5 was used for binding of the enzyme and the color eluted the pure enzyme was eluted using a NaCl gradient.

Characterization:

The purified protein gave a single band in SDS-PAGE of 50
20 kDa with a pI of around 3.7.

The specific activity on ECU was 50 ECU per mg of protein. A molar extinction value of 113380 was based on the amino acid composition and was used for determination of the protein concentration after a ratio of 1.8 (E280/E260) was obtained and us-
25 ing the MW of 50 kDa.

Catalytic activity on phosphoric acid swollen cellulose at pH 8.5 was apparent k_{cat} of 22 per sec and apparent K_M of 0.6 g/l. The cellulase has no activity on PNP-beta cellobioside.

Buffers

30 All enzyme experiments were performed in one of the following buffers:

pH 3.0 to 3.5:

0.1 M sodium citrate

pH 4.0 to 5.5:	0.1 M sodium acetate
pH 6.0:	0.1 M sodium MES buffer
pH 6.5 to 7.5:	0.1 M sodium MOPS buffer
pH 8.0 to 8.5:	0.1 M Barbitol buffer
5 pH 9.0 to 10.0:	0.1 M Glycine buffer

pH activity profiles

The pH activity profiles were obtained using CMC. The final CMC concentration was 7.5 gram per liter, incubation was for 20 min at 40°C and the formation of reducing sugars determined by
10 using *p*-hydroxy-benzoic-acid-hydrazide (PHBAH) modified from Lever (1972) using 5 g of potassium sodium tartrate in addition to 1.5 g of PHBAH. The pH optimum was 8.0, and more than 50% relative activity was obtained between pH 6.5 and 9.0.

Apparent kinetic constant determination using phosphoric-acid 15 swollen cellulose (PASC)

PASC stock solution was prepared as follows: 5 g of cellulose (Avicel) was moistened with water, and 150 ml ice cold 85% ortho-phosphoric-acid was added. The suspension was slowly stirred in an ice-bath for 1 h. Then 100 ml ice cold acetone was
20 added while stirring. The slurry was transferred to a Buchner filter with Pyrex sintered disc number 3 and then washed three times with 100 ml ice cold acetone, sucked as dry as possible after each wash. Finally it was washed twice with 500 ml water, and again sucked as dry as possible after each wash. The PASC
25 was mixed with deionized water to a total volume of 300 ml. It was blended to homogeneity (using an Ultra Turrax Homogenizer) and stored in a refrigerator for up to one month.

Substrate was equilibrated with buffer using the following procedure: 20 g phosphoric-acid swollen cellulose PASC stock so-
30 lution was centrifuged for 20 min at 5000 rpm, the supernatant was poured off, and the sediment was resuspended in 30 ml of buffer. After 20 min centrifugation at 5000 rpm, the supernatant was decanted, and the sediment was resuspended in buffer to a

total of 30 g. This corresponds to a substrate concentration of 10 mg l⁻¹.

To measure kinetic parameters, substrate concentrations from 0.2 mg ml⁻¹ to 8 mg ml⁻¹ were used. Rates were measured at 8 different substrate concentrations in duplicate. The amount of reducing sugars was determined using the PHBAH method modified from Lever (1972).

The enzyme concentration was calculated using the molar absorbancy. The apparent kinetic constants $K_M(\text{app.})$, $V_{\text{max}}(\text{app.})$ and $k_{\text{cat}}(\text{app.})$ were calculated using the equation for enzyme kinetics in the computer program GraFit (Leatherbarrow, 1992).

EXAMPLE 4

Tensile strength loss induced by treating fabric with endo- β -1,4-glucanase from *Saccharothrix australiensis*, IFO 14444

Tensile strength loss induced by treating textile with the purified endoglucanase from example 3 was measured and compared with a fungal family 6 endoglucanase.

Experimental protocol:

- | | | |
|----|-------------|---|
| 20 | Buffer | : 0.05 M phosphate buffer pH 7.0 |
| | Volume | : 100 ml |
| | Temperature | : room temperature |
| | Textile | : Tea towels pre-aged ED 9613829 5 pieces á 5 x 25 cm |
| 25 | Dosage | : 2x0, 1000, 10.000, 100.000 ECU/l Carezyme 59100 ECU/l MB145
1900, 19.000, 190.000 ECU/l of family 6 cellulases |
| | Time | : 7 days dark storage |
| 30 | Rinse time | : 10 min. in running tap water |
| | Evaluation | : Tensile strength loss measured on an Instron instrument 5564. Breakage within 20+/-3 sec. |

Pre-aged fabric was incubated for a week with different dosages of enzyme in phosphate buffer pH 7.0. Afterwards the

fabric was rinsed in water, acclimatised in climate room for 24 hours (60 % RH, 20°C). Tensile strength was measured on an Instron instrument. The tensile strength loss is calculated relative to pre-aged textile which has not been incubated (reference). Blank (no enzyme) and the EG VI from the fungus *Humicola insolens* (family 6 endoglucanase) were included in the experiment for comparison.

Results:

	ECU/l	% TSL
Reference		0
Blank	0	-3.6
Fungal EG VI	19,000	1.2
	190,000	12.5
Enzyme of the invention	59,100	-6.7

The enzyme of the invention gave not rise to any tensile strength loss.

EXAMPLE 5**Colour Clarification in Terg-O-Meter**

In this example the capability of the endoglucanase of the invention to rejuvenate the colour of cotton textile is demonstrated using an assay for determining colour care benefits, i.e. "Color Clarification", of cotton cloth in a miniaturised washing machine, the 100 ml Terg-O-Meter.

250 ml beakers with 100 ml buffer (or detergent) was positioned in a Terg-O-Meter and equilibrated to 35°C. Then two 7x7 cm swatches of black, woven cotton cloth was added to each beaker, the stirrers were put in motion, and finally enzyme was added: A) A blank, B) Three different dosages of a standard (e.g. the commercial available enzyme preparation Celluzyme™), and C) Two different dosages of the inventive endoglucanase. Incubation then proceeded for 30 minutes at 35°C

After the 30 minutes of incubation the swatches were rinsed in cold tap water for 10 minutes and dried in a tumble dryer.

The cycle of incubation and rinsing/drying was repeated once - or until the swatches clearly differed in respect to colour and/or fuzz in the swatches surface.

Finally the swatches were graded against the blank (no enzyme) and the standard (e.g. Celluzyme) swatches. Visual grading was performed by a panel of trained graders, and colour was measured with a remission spectrometer. Results are expressed in Table 1 as "Colour Clarification" (CC) of the swatches obtained per activity unit of enzyme.

10

Table 1:

Enzyme	Cellulase Family	CC - black, woven
Celluzyme	multi-component	1.0
Blank	no enzyme	0.0
Saccharothrix australiensis	endoglucanase	1.8

LITERATURE

Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995.

Axelsen, N., et al. in: A Manual of Quantitative Immuno-electrophoresis, Blackwell Scientific Publications, 1973, Chapter 23.

Denman, S. et al.: Characterization of a *Neocallumastix patriarum* cellulase cDNA (*celA*) homologous to *Trichoderma reesei* cellobiohydrolase II, Appl. Environ. Microbiol. (1996), 62(6), 1889-1896.

Damude, H.G. et al.: Substrate specificity of endoglucanase A from *Cellulomonas fimi*: fundamental differences between endoglucanases and exoglucanases from family 6, Biochem.J. (1996), 315(2), 467-72.

Devereux et al. (1984) Nucleic Acids Res. 12, 387-395.

Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of *aldB*, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*. J. Bacteriol., 172, 4315-4321

Dretzen, G., Bellard, M., Sassone-Corsi, P., Chambon, P. (1981) A reliable method for the recovery of DNA fragments from agarose and acrylamide gels. Anal. Biochem., 112, 295-298.

Feinberg, A. P. and Vogelstein, B. (1983) Anal. Biochem. 132:6-13.

Gilbert, H.J. and Hazlewood, G.P. (1993) J. Gen. Microbiol. 139:187-194.

Gilkes, N.R., Henrissat, B., Kilburn, D.G., Miller Jr., R.C. and Warren, R.A.J.: Domains in microbial β -1,4-glycanases; sequence

conservation, function, and enzyme families. Microbiol. Rev. 55 (1991), 305-315.

Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990.

Henrissat, B.: A classification of glycosyl hydrolases based on amino acid sequence similarities. Biochem. J. 280 (1991), 309-316.

Henrissat, B., and Bairoch, A.: New families in the classification of glycosyl hydrolases based on amino acid sequence similarities. Biochem. J. 293 (1993), 781-788.

Johnstone, A. and R. Thorpe, Immunochemistry in Practice, Blackwell Scientific Publications, 1982 (pp. 27-31).

Leatherbarrow, R. J. (1992) Grafit version 3.0 Erithacus Software Ltd. Staines, U.K.

- 5 Lever, M. (1972) A new reaction for colormetric determination of carbohydrates. Anal. Biochem. 47, 273-279.

O. Ouchterlony in: Handbook of Experimental Immunology (D.M. Weir, Ed.), Blackwell Scientific Publications, 1967, pp. 655-706.

Pitcher, D. G., Saunders, N. A., Owen, R. J. (1989): Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. Lett. Appl. Microbiol., 8, 151-156).

Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY.

- 10 Quillet, L. et al.: The gene encoding the beta-1,4-endoglucanase (Cel) from *Myxococcus xanthus*: evidence for independent acquisition by horizontal transfer of binding and catalytic domains from actinomycetes, Gene (1995), 158(1), 23-9.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT:
 (A) NAME: NOVO NORDISK A/S
 (B) STREET: Novo Alle
 (C) CITY: Bagsvaerd
10 (E) COUNTRY: Denmark
 (F) POSTAL CODE (ZIP): DK-2880
 (G) TELEPHONE: +45 44 44 88 88
 (H) TELEFAX: +45 44 49 32 56
- 15 (ii) TITLE OF INVENTION: ENDO-BETA-1,4-GLUCANASES FROM SAC-
CHAROTHRIX
- (iii) NUMBER OF SEQUENCES: 2
- 20 (iv) COMPUTER READABLE FORM:
 (A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
25 (EPO) (D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(2) INFORMATION FOR SEQ ID NO: 1:

- 30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1470 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:
- | | | |
|----|--|-----|
| 40 | ATGCACCCCC GCTCGAAGAG ACCCCTCACC ACCAGACGCA AGGTCGTCCC GGCCGTCGCG | 60 |
| | GCCGGAACCG TCCTCGCCGG CGGCGTCACC GCCCTGACCT CCAACATCGC GCAGGCCGCC | 120 |
| | GCCGGCTGCC GCGTCGACTA CGCCGTGACG AGCCAGTGGC CCGGTGGCTT CGGTGCAGCC | 180 |
| 45 | GTCACCGTCA CGAACCTCGG CGACCCGCTC TCGTCCTGGG AGCTGAGCTG GACGTTCCCC | 240 |
| | GACGGCCAGG GCGTGCAGCA GCTCTGGAAC GGCCTGCACT CGACCTCCGG TTCGAACGTC | 300 |
| 50 | ACCGTGAAAG AAATGTCGTG GAACGGTTCG GTCGGCACCA ACGCCAGCGT CCAGGTCGGC | 360 |
| | TTCAACGGCT CCTGGAACGG CGCGAACAAC GCGCCGACGT CCTTCACGCT CAACGGCACC | 420 |
| | TCTGTCAACG GTGCGGTCGG TGGCCCCGACG ACGGAGCCGA CGCCCGAGCC GACCCCGGAG | 480 |
| 55 | CCCACGCCCC AGCCGACGCC GGAGCCGACG CCCGAGCCGA CGCCGAGGCC CACGCCCGAG | 540 |

CCGACGCCGG AGCCACGCC CGAGCCGACC CCGGAGCCCA CGCCCGAGCC CACGCCCGAG 600
 CCCACGCCCC AGCCACGAT GCCGCCGGTC CAGGCCGGTC AGTTCCACGT CGACACCACG 660
 5 AACCAGTCGT ACCGCGCCTG GCAGGCGGCC AGCGGCTCCG ACAAGGACCT GCTGGCGAAG 720
 ATCGCCCTGA CGCCGAGGC GTACTGGGTC GGCAACTGGA ACGAAGCCTC GCACGCGCAG 780
 CAGGAAGTCC GTGACATCAC GTCGGCCGCT GCGGCCGCCG GCAGGACCGC CGTGCTCGTC 840
 10 GTCTACGCCA TCCCGGGCCG CGACTGCGGC CAGCACTCCA GCGGCGGCGT GTCGACCTCC 900
 GAGTACGCGC AGTGGATCGA CACGGTCGCC CAGGGCATCG TCGGCAACCC GTGGGTGGTC 960
 15 CTCGACCCCG ACGCGTGCC GATGCTCGGC GACTGCGACG GCCAGGGCGA CCGGGTCGGC 1020
 TTCCTCAAGT ACGCCGCGAA GTCCCTGACC GCCAAGGGTG CGCGGTCTA CATCGACGCC 1080
 GGCCACTCGG CGTGGCTGTC GCCGTCGAA GCCGCGAACC GCCTCAACCA GATCGGGTTC 1140
 20 GAGGACGCCG TGGGCTTCTC GATCAACGTC TCCAACCTACC GCACGACGGC GGAGTCGAAG 1200
 ACCTGGGGTC AGCAGGTCTC GCAGCTGACC GGTGGCAAGA AGTTCGTCAT CGACACGTCG 1260
 25 CGCAACGGCA ACGGCCCGTC CGGGTCGAA TGGTGCAACC CGAGCGGCCG CGCCCTCGGC 1320
 GAGCGCCCGA CGCTCGTGAA CGACCGCAGC GGGCTCGACG CGCTGCTGTG GATCAAGCTG 1380
 CCCGGTGAGT CGGACGGCGC CTGCAACGGC GGCCCGGGCG CCGGTCAGTG GTGGCACTCC 1440
 30 ATGGCACTGG CTGGCCGCAA CGCGAAGTGG 1470

(
 2) INFORMATION FOR SEQ ID NO: 2:

35

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 490 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

45

Met His Pro Arg Ser Lys Arg Pro Leu Thr Thr Arg Arg Lys Val Val
 1 5 10 15
 Pro Ala Val Ala Ala Gly Thr Val Leu Ala Gly Gly Val Thr Ala Leu
 50 20 25 30
 Thr Ser Asn Ile Ala Gln Ala Ala Ala Gly Cys Arg Val Asp Tyr Ala
 35 40 45
 55 Val Thr Ser Gln Trp Pro Gly Gly Phe Gly Ala Ala Val Thr Val Thr
 50 55 60

70

	Asn	Leu	Gly	Asp	Pro	Leu	Ser	Ser	Trp	Glu	Leu	Ser	Trp	Thr	Phe	Pro	
	65					70					75					80	
5	Asp	Gly	Gln	Gly	Val	Gln	Gln	Leu	Trp	Asn	Gly	Val	His	Ser	Thr	Ser	
					85					90					95		
	Gly	Ser	Asn	Val	Thr	Val	Lys	Glu	Met	Ser	Trp	Asn	Gly	Ser	Val	Gly	
				100					105					110			
10	Thr	Asn	Ala	Ser	Val	Gln	Val	Gly	Phe	Asn	Gly	Ser	Trp	Asn	Gly	Ala	
			115					120					125				
	Asn	Asn	Ala	Pro	Thr	Ser	Phe	Thr	Leu	Asn	Gly	Thr	Ser	Cys	Asn	Gly	
			130				135					140					
15	Ala	Val	Gly	Gly	Pro	Thr	Thr	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	
	145					150					155					160	
	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	
20					165				170						175		
	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	
				180				185					190				
25	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Pro	Glu	Pro	Thr	Met	Pro	
			195				200					205					
	Pro	Val	Gln	Ala	Gly	Gln	Phe	His	Val	Asp	Thr	Thr	Asn	Gln	Ser	Tyr	
		210					215					220					
30	Arg	Ala	Trp	Gln	Ala	Ala	Ser	Gly	Ser	Asp	Lys	Asp	Leu	Leu	Ala	Lys	
	225				230						235					240	
	Ile	Ala	Leu	Thr	Pro	Gln	Ala	Tyr	Trp	Val	Gly	Asn	Trp	Asn	Glu	Ala	
35					245					250					255		
	Ser	His	Ala	Gln	Gln	Glu	Val	Arg	Asp	Ile	Thr	Ser	Ala	Ala	Ala	Ala	
				260					265					270			
40	Ala	Gly	Arg	Thr	Ala	Val	Leu	Val	Val	Tyr	Ala	Ile	Pro	Gly	Arg	Asp	
			275				280						285				
	Cys	Gly	Gln	His	Ser	Ser	Gly	Gly	Val	Ser	Thr	Ser	Glu	Tyr	Ala	Gln	
		290					295					300					
45	Trp	Ile	Asp	Thr	Val	Ala	Gln	Gly	Ile	Val	Gly	Asn	Pro	Trp	Val	Val	
	305					310					315					320	
	Leu	Asp	Pro	Asp	Ala	Leu	Pro	Met	Leu	Gly	Asp	Cys	Asp	Gly	Gln	Gly	
50					325					330					335		
	Asp	Arg	Val	Gly	Phe	Leu	Lys	Tyr	Ala	Ala	Lys	Ser	Leu	Thr	Ala	Lys	
				340					345					350			
55	Gly	Ala	Arg	Val	Tyr	Ile	Asp	Ala	Gly	His	Ser	Ala	Trp	Leu	Ser	Pro	
			355				360						365				
	Ser	Glu	Ala	Ala	Asn	Arg	Leu	Asn	Gln	Ile	Gly	Phe	Glu	Asp	Ala	Val	

71

	370		375		380	
	Gly Phe Ser Ile Asn Val Ser Asn Tyr Arg Thr Thr Ala Glu Ser Lys					
	385		390		395	400
5	Thr Trp Gly Gln Gln Val Ser Gln Leu Thr Gly Gly Lys Lys Phe Val					
			405		410	415
	Ile Asp Thr Ser Arg Asn Gly Asn Gly Pro Ser Gly Ser Glu Trp Cys					
10		420		425		430
	Asn Pro Ser Gly Arg Ala Leu Gly Glu Arg Pro Thr Leu Val Asn Asp					
		435		440		445
15	Arg Ser Gly Leu Asp Ala Leu Leu Trp Ile Lys Leu Pro Gly Glu Ser					
		450		455		460
	Asp Gly Ala Cys Asn Gly Gly Pro Gly Ala Gly Gln Trp Trp His Ser					
	465		470		475	480
20	Met Ala Leu Ala Gly Arg Asn Ala Lys Trp					
		485		490		


Applicant's or agent's file reference number	5195-WO, AMJ	International application No.	PCT/DK 98 / 00286
---	--------------	-------------------------------	-------------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

5

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>4</u> , lines <u>13-21</u> .	
B. IDENTIFICATION OF DEPOSIT	Further deposits are identified on an additional sheet <input type="checkbox"/>
Name of depositary institution DEUTSCHE SAMMLUNG VON MIKROORGANISMEN UND ZELLKULTUREN GmbH	
Address of depositary institution (including postal code and country) Mascheroder Weg 1b, D-38124 Braunschweig, GERMANY	
Date of deposit 17 March 1997	Accession Number DSM 11476
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional <input type="checkbox"/>	
Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input checked="" type="checkbox"/>	This sheet was received with the international application
Authorized officer 	

For International Bureau use only	
<input type="checkbox"/>	This sheet was received by the International Bureau on:
Authorized officer	

CLAIMS

1. An enzyme preparation comprising an enzyme having endo- β -1,4-glucanase activity which enzyme is obtainable from or
5 endogeneous to a strain belonging to the genus *Saccharothrix*.

2. The preparation according to claim 1, wherein the strain is selected from the group of species consisting of *Saccharothrix australiensis*, *Saccharothrix texasensis*, *Saccharothrix*
10 *waywayandensis*, *Saccharothrix cryophilis*, *Saccharothrix flava*, *Saccharothrix coeruleofusca*, *Saccharothrix longispora*, *Saccharothrix mutabilis* ssp. *capreolus*, *Saccharothrix aerocolonigenes*, *Saccharothrix mutabilis* ssp. *mutabilis*, *Saccharothrix syringae*, *Saccharothrix* sp.

15

3. The preparation according to claim 2, wherein the strain is the species *Saccharothrix australiensis*, IFO 14444, *Saccharothrix texasensis*, NRRL B-16134, *Saccharothrix*
waywayandensis, NRRL B-16159, *Saccharothrix cryophilis*, NRRL B-
20 16238, *Saccharothrix flava*, ATCC 29533, *Saccharothrix* *coeruleofusca*, ATCC 35108, *Saccharothrix longispora*, ATCC 31109, *Saccharothrix mutabilis* ssp. *capreolus*, ATCC 23892, *Saccharothrix aerocolonigenes*, ATCC 23870, *Saccharothrix mutabilis* ssp. *mutabilis*, ATCC 31520, *Saccharothrix syringae*,
25 DSM 43886, or *Saccharothrix* sp., IFO 13785.

30

4. The preparation according to any of the claims 1-3, wherein the endo- β -1,4-glucanase belongs to family 6 of glycosyl hydrolases.

5. The preparation according to any of the claims 1-4, wherein the endo- β -1,4-glucanase is active at a pH in the range of 4-11, preferably 5.5-10.5.

35 6. An endo- β -1,4-glucanase which is

(a) a polypeptide produced by *Saccharothrix australiensis*, IFO 14444, or

- (b) a polypeptide comprising an amino acid sequence as shown in positions 226-490 of SEQ ID NO:2, or
- (c) an analogue of the polypeptide defined in (a) or (b) which is at least 75% homologous with said polypeptide, or is derived from said polypeptide by substitution, deletion or addition of one or several amino acids, or is immunologically reactive with a polyclonal antibody raised against said polypeptide in purified form.
- 10 7. An isolated polynucleotide molecule encoding a polypeptide having endo-beta-1,4-endoglucanase activity selected from the group consisting of:
- (a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 676 to nucleotide 1470;
- 15 (b) polynucleotide molecules that encode a polypeptide that is at least 80% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 226 to amino acid residue 490; and
- (c) degenerate nucleotide sequences of (a) or (b).
- 20 8. The isolated polynucleotide molecule according to claim 7, wherein the polynucleotide is DNA.
9. An isolated polynucleotide molecule encoding a polypeptide having endo-beta-1,4-glucanase activity which polynucleotide
- 25 molecule hybridizes to a denatured double-stranded DNA probe under medium stringency conditions, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 676-1470 of SEQ ID NO:1 and DNA probes comprising a subsequence of positions 676-1470 of SEQ ID NO:1 having a length of at least about 100 base pairs.
- 30 10. The isolated polynucleotide molecule according to claim 9 which is isolated from or produced on the basis of a DNA library from a prokaryote, preferably from a bacterium, more preferably from a gram positive bacterium.

11. The isolated polynucleotide molecule according to claim 10 which is isolated from or produced on the basis of a DNA library from a strain belonging to the class Actinomycetes, preferably
5 to the family Pseudonocardiaceae, more preferably to the genus *Saccharothrix*, in particular a strain of *Saccharothrix australiensis*, especially *Saccharothrix australiensis*, IFO 14444.

10 12. The isolated polynucleotide molecule according to any of the claims 7-11 which is isolated from *Escherichia coli*, DSM 11476.

13. The isolated polynucleotide molecule according to any of the claims 7-12 which further comprises a partial DNA sequence
15 encoding a cellulose binding domain (CBD).

14. The isolated polynucleotide molecule according to claim 13, wherein the partial nucleotide sequence encoding a cellulose binding domain (CBD) corresponds to the nucleotides in positions
20 60-435 of SEQ ID NO:1.

15. The isolated polynucleotide molecule according to claim 13 or 14, which further comprises a partial nucleotide sequence encoding a linking region, the linking region operably linking
25 the cellulose binding domain (CBD) and the catalytically active domain (CAD) of the enzyme encoded by the nucleotide sequence comprised by the isolated polynucleotide molecule.

16. An expression vector comprising the following operably
30 linked elements: a transcription promoter; a DNA segment selected from the group consisting of (a) polynucleotide molecules encoding a polypeptide having endo-beta-1,4-glucanase activity comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 676 to nucleotide 1470, (b) polynucleotide molecules
35 encoding a polypeptide having endo-beta-1,4-glucanase activity that is at least 75% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 226 to amino acid residue 490,

and (c) degenerate nucleotide sequences of (a) or (b); and a transcription terminator.

17. A cultured cell into which has been introduced an expression
5 vector according to claim 16, wherein said cell expresses the polypeptide encoded by the DNA segment.

18. The cell according to claim 17, which is a prokaryotic cell,
in particular a bacterial cell, or an endogenous cell from which
10 the DNA segment, encoding the polypeptide exhibiting endoglucanase activity, originates.

19. The cell according to claim 17, wherein the cell belongs to
a strain of *Bacillus*, preferably a strain of *Bacillus subtilis*
15 or *Bacillus lentus*.

20. A cell according to claim 17, wherein the cell belongs to a
strain of *Saccharothrix australiensis*, preferably *Saccharothrix*
australiensis, IFO 14444.

20

21. The cell according to claim 17, wherein the cell belongs to
a strain of *Pseudomonas*, preferably a strain of *Pseudomonas*
fluorescens or *Pseudomonas mendocina*.

22. The cell according to claim 17, wherein the cell belongs to
25 a strain of *Streptomyces*.

23. A cell according to claim 17 wherein the cell belongs to a
strain of *Saccharomyces*, preferably a strain of *Saccharomyces*
30 *cerevisiae*.

24. A method of producing a polypeptide having endo-beta-1,4-
glucanase activity comprising culturing a cell into which has
been introduced an expression vector according to claim 16,
35 whereby said cell expresses a polypeptide encoded by the DNA
segment; and recovering the polypeptide.

25. An enzyme preparation comprising a purified polypeptide according to claim 6.
26. The preparation according to claim 1 or 25 which further
5 comprises one or more enzymes selected from the group consisting of proteases, cellulases (endoglucanases), β -glucanases, hemicellulases, lipases, peroxidases, laccases, α -amylases, glucoamylases, cutinases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, pectate lyases, xyloglu-
10 canases, xylanases, pectin acetyl esterases, polygalacturonases, rhamnogalacturonases, pectin lyases, other mannanases, pectin methylesterases, cellobiohydrolases, transglutaminases; or mixtures thereof.
- 15 27. An isolated enzyme having endo-beta-1,4-glucanase activity, in which the enzyme is (i) free from homologous impurities, and (ii) produced by the method according to claim 24.
28. An isolated substantially pure biological culture of the
20 strain *Escherichia coli*, DSM 11476.
29. Use of the enzyme according to claim 6 or 27 or the enzyme preparation according to any of the claims 1-5, 25 and 26 in the textile industry for improving the properties of cellulosic
25 fibres or fabric or for providing a stone-washed look of denim; or in industrial cleaning processes.
30. A detergent composition comprising the enzyme according to claim 6 or 27 or the enzyme preparation according to any of the
30 claims 1-5, 25 and 26.
31. A fabric softener composition comprising the enzyme according to claim 6 or 27 or the enzyme preparation according to any of the claims 1-5, 25 and 26.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00286

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C12N 9/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, BIOSIS, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Applied and Environmental Microbiology, Volume 54, No 10, October 1988, Gurdev S. Ghangas et al, "Cloning of the Thermomonospora fusca Endoglucanase E2 Gene in Streptomyces lividans: Affinity Purification and Functional Domains of the Cloned Gene Product" page 2521 - page 2526 --	1-31
X	US 5643791 A (R. ANTHONY J. WARREN ET AL), 1 July 1997 (01.07.97), figure 5, abstract --	6,9,16-18, 24-27,29-31

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

9 October 1998

Date of mailing of the international search report

13-10- 1998

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Hampus Rystedt

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00286

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>Canadian journal of microbiology, Volume 40, No 5, 1994, Ethier, J.-F. et al, "Cloning of two xylanase genes from the newly isolated actinomycete Actinomadura sp. strain FC7 and characterization of the gene products" page 362 - page 368</p> <p style="text-align: center;">-- -----</p>	1-31

Information on patent family members

International application No.

PCT/DK 98/00286

Form PCT/ISA/210 (patent family annex) (July 1992)